

# The Experiences of Young Adults With Hodgkin Lymphoma Transitioning to Survivorship: A Grounded Theory Study

Lauren Matheson, PhD, MSc, BSc, Mary Boulton, PhD, BA (Hons),  
Verna Lavender, PhD, PGCE, BSc (Hons), RGN, Graham Collins, MD, DPhil,  
Tracy Mitchell-Floyd, BA (Hons), RN, and Eila Watson, PhD, BSc (Hons)

Matheson is a postdoctoral research assistant, Boulton is a professor of health sociology, and Lavender is a senior lecturer in cancer care, all in the Faculty of Health and Life Sciences at Oxford Brookes University; Collins is a consultant hematologist and Mitchell-Floyd is a lymphoma specialist nurse practitioner, both in the Department of Clinical Haematology at the Oxford Cancer and Haematology Centre of the Oxford University Hospitals NHS Foundation Trust; and Watson is a professor of supportive cancer care in the Faculty of Health and Life Sciences at Oxford Brookes University, all in Oxford, England.

Matheson, Boulton, Lavender, and Watson contributed to the conceptualization and design and provided the analysis. Matheson completed the data collection, and Collins and Mitchell-Floyd assisted with recruitment. All authors contributed to the manuscript preparation.

Matheson can be reached at [l.matheson@brookes.ac.uk](mailto:l.matheson@brookes.ac.uk), with copy to editor at [ONFEditor@ons.org](mailto:ONFEditor@ons.org).

Submitted November 2015. Accepted for publication January 21, 2016.

Key words: young adults; cancer survivors; qualitative; grounded theory; Hodgkin lymphoma

ONF, 43(5), E195-E204.

doi: 10.1188/16.ONF.E195-E2014

**Purpose/Objectives:** To explore the experiences of young adults with Hodgkin lymphoma during the first year following the end of initial treatment.

**Research Approach:** A qualitative grounded theory study.

**Setting:** Interviews with patients recruited from three cancer centers in England.

**Participants:** 10 Hodgkin lymphoma survivors (four men and six women aged 21–39 years) recruited as part of a larger study of 28 young adult cancer survivors.

**Methodologic Approach:** Semistructured interviews were conducted about two months after treatment completion, and follow-up interviews were conducted seven months later. The authors' grounded theory of positive psychosocial adjustment to cancer provided the conceptual framework.

**Findings:** Positive reframing, informal peer support, acceptance, and normalization helped young adults dismantle the threats of Hodgkin lymphoma during the course of treatment. However, they described losing a sense of security following treatment completion. Greater age-specific information to enable better preparation for the future was desired regarding body image, fertility, sexual relationships, work, and socializing.

**Conclusions:** Informal support mechanisms, like peer support and patient navigator interventions, may be useful ways to further support young adults after treatment completion.

**Interpretation:** Positive psychosocial adjustment to cancer survivorship in young adults is facilitated by having informal peer support; being able to positively reframe, accept, and normalize their experience; and being prepared for the future.

Young adult cancer survivors have unique experiences, needs, and challenges dependent on their age and diagnosis (Hall et al., 2012; Zebrack, 2009; Zebrack & Isaacson, 2012). Cancer can disrupt normal life stages, such as finding a partner, establishing independence, finishing education, starting a family, and gaining employment (Grinyer, 2009; Zebrack, 2011). Young adult cancer survivors may also have specific concerns and needs regarding appearance and body image, identity development, and long-term future health (Bellizzi et al., 2012; Kelly & Gibson, 2008; Zebrack, 2011; Zebrack & Isaacson, 2012). Greater levels of psychological distress have been shown in young adult survivors compared to older survivors (Costanzo et al., 2007; Costanzo, Ryff, & Singer, 2009; Kwak et al., 2013; Parker, Baile, de Moor, & Cohen, 2003); therefore, young adults require specific interventions from the oncology nurse (Adams et al., 2011; Hall et al., 2012; Zebrack, 2009; Zebrack & Isaacson, 2012).

The transition from initial treatment completion—commonly a course of 4–6 monthly cycles of cytotoxic chemotherapy treatment—to follow-up care can be