

# The Effectiveness of Yoga on Cancer-Related Fatigue: A Systematic Review and Meta-Analysis

Jihong Song, MA, Tao Wang, MA, Yujie Wang, MA, Rong Li, MA, Sitian Niu, BSN, Litao Zhuo, BSN, Qian Guo, MA, and Xiaomei Li, PhD

**PROBLEM IDENTIFICATION:** The aim of this article is to evaluate the effectiveness of yoga on cancer-related fatigue (CRF) in patients undergoing chemotherapy and/or radiation therapy.

**LITERATURE SEARCH:** Relevant English and Chinese articles were retrieved from medical databases and included in this analysis. Standardized critical appraisal instruments from the Joanna Briggs Institute were adopted for the quality assessment.

**DATA EVALUATION:** 16 randomized controlled trials met the inclusion criteria.

**SYNTHESIS:** Yoga interventions had a positive effect in reducing CRF among patients undergoing chemotherapy and/or radiation therapy, but the adherence to yoga was low. Mixed types of yoga, in addition to supervised and self-practicing strategies, were associated with increased patient adherence and improved CRF.

**IMPLICATIONS FOR PRACTICE:** Yoga appears to be a safe and effective exercise for the management of CRF during chemotherapy and/or radiation therapy; however, additional high-quality studies are needed to define an optimal yoga intervention strategy.

**KEYWORDS** yoga; fatigue; chemotherapy; radiation therapy; meta-analysis; systematic review

**ONF, 48(2), 207-228.**

**DOI** 10.1188/21.ONF.207-228

Cancer is the second leading cause of death in the world, with an estimated 9.6 million deaths occurring in 2018 (World Health Organization, 2018). According to a cancer progress report released by the American Association for Cancer Research (2018), the number of new cancer cases worldwide could increase to 24 million annually by 2035. With advancements in cancer prevention, diagnosis, and treatment, survival rates are continually increasing; however, the physical and psychological disorders associated with the occurrence, progression, and treatment of cancer, including nausea, poor appetite, fatigue, decreased immunity, anxiety, and depression, significantly affect the quality of life of cancer survivors.

Cancer-related fatigue (CRF) is one of the most common symptoms associated with cancer and is defined as “a distressing, persistent, subjective sense of physical, emotional, and/or cognitive tiredness or exhaustion related to cancer or cancer treatment that is not proportional to recent activity and interferes with the usual function” (National Comprehensive Cancer Network, 2018, p. MS-3). Research has shown that 60% to 99% of patients undergoing chemotherapy and/or radiation therapy have reported CRF (Wu et al., 2017). Exercise has been examined as one way to reduce CRF. The American College of Sports Medicine has recommended that an overall volume of weekly activity consisting of moderate-intensity exercise for 150 minutes, vigorous-intensity exercise for 75 minutes, or an equivalent combination is appropriate for cancer survivors (Schmitz et al., 2010). However, for patients undergoing chemotherapy and/or radiation therapy, several reviews have reported that shorter exercise sessions or lower targeted exercise volume were associated with greater improvements in CRF (Carayol et al., 2015; Kessels et al., 2018; Tian et al., 2016).

Yoga has been identified as one such low-intensity exercise. Yoga has its roots in ancient Indian philosophy and, differing from general aerobic exercise, yoga requires that the practitioner focuses on the integration, unity, and harmony of inner awareness, breathing, and body (Carayol et al., 2013; Cramer et al., 2017). There are various styles of yoga related to different yoga schools. Therapeutic yoga for patients with cancer commonly involves one or more physical postures, breathing techniques, and meditation. Research has demonstrated that yoga can modulate hypothalamic-pituitary-adrenal dysregulation and improve mental and physical health (Streeter et al., 2012). However, to date, insufficient evidence exists to substantiate the effectiveness of yoga on CRF in patients undergoing chemotherapy and/or radiation therapy.

A Cochrane review by Cramer et al. (2017) of 24 randomized controlled trials (RCTs), with a total

of 2,166 women undergoing active treatment for breast cancer, found that yoga reduced CRF when compared to no therapy or psychoeducation only. However, the review only included patients with breast cancer and subgroup analysis could not be performed. In contrast, a meta-analysis by Pan et al. (2015) of nine RCTs revealed no significant relief of fatigue from yoga; however, data concerning CRF during treatment were not extracted independently.


The effectiveness of a yoga intervention as well as the amount of yoga to be performed by patients with CRF remains in question. Several reviews have attempted to examine this topic, but most have not been able to perform a meta-analysis, mostly related to small sample sizes and limited database searching (Danhauer, Addington, et al., 2015; Danhauer et al., 2019; Felbel et al., 2014; Harder et al., 2012; Pan et al., 2015; Sadjja & Mills, 2013; Smith & Pukall, 2009; Tolia et al., 2018; Zuo et al., 2016); several have focused only on one cancer, particularly breast cancer (Felbel et al., 2014; Harder et al., 2012; Pan et al., 2015; Zuo et al., 2016); and some have not extracted and analyzed information about CRF in patients during chemotherapy and/or radiation therapy (Buffart et al., 2012; Carayol et al., 2015; Pan et al., 2015; Tomlinson et al., 2014). Therefore, additional large-scale data examinations are needed to determine the effectiveness of yoga on CRF in patients undergoing chemotherapy and/or radiation therapy.

Several clinical trials have been conducted to investigate the effectiveness of yoga on CRF among cohorts of patients treated for different cancers (Ben-Josef et al., 2017; Cramer et al., 2016; Lötze et al., 2016). The aim of the current review was to assess the effectiveness and appropriate amount of yoga as an intervention for adults experiencing CRF while undergoing chemotherapy and/or radiation therapy, compared with no therapy or psychoeducation. In addition, the optimal yoga intervention strategy and style for reducing CRF, as well as patient adherence and factors affecting patient adherence, were investigated. The safety and adverse events of a yoga intervention during chemotherapy and/or radiation therapy were also examined.

## Methods

The systematic review was performed according to an established protocol (PROSPERO No. CRD42020156435) and presented using the Preferred Reporting Items for Systematic Reviews and Meta-Analyses guidelines (Moher et al., 2015).

**FIGURE 1. PRISMA Flow Diagram of Literature Search**



**TABLE 1. Characteristics of Included Studies (N = 16)**

Study and Location	Sample	Intervention	Duration	Endpoint
Ben-Josef et al., 2017 (United States)	Male patients with prostate cancer undergoing radiation therapy; 35 were enrolled in an intervention group and 33 in a control group.	Eischens yoga (mixed); supervised	75 minutes per day, 2 days per week	6–9 weeks
Chakrabarty et al., 2015 (India)	Female patients with breast cancer undergoing radiation therapy; 80 were enrolled in an intervention group and 80 in a control group.	Pranayama (mind-breathing); supervised	18 minutes twice per day, 5 days per week	6 weeks
Chandwani et al., 2010 (United States)	Female patients with stage 0–III breast cancer undergoing radiation therapy; 30 were enrolled in the intervention group (27 completed), and 31 were enrolled in the control group (29 completed).	Patanjali yoga tradition; mixed plus CD	60 minutes per day, 3 days per week	6 weeks
Chandwani et al., 2014 (United States)	Female patients with stage 0–III breast cancer undergoing radiation therapy; 53 were enrolled in a yoga intervention (49 completed), and 56 were enrolled in a stretching intervention (52 completed). 54 were enrolled in a control group (48 completed).	Asanas plus pranayama yoga; mixed plus CD	60 minutes per day, 3 days per week	6 weeks
Chaoul et al., 2018 (United States)	Female patients with breast cancer undergoing chemotherapy; 74 were enrolled in a yoga intervention (64 completed), and 68 were enrolled in a stretching intervention (59 completed). 85 were enrolled in a control group (79 completed).	Tibetan yoga (mind-breathing); mixed plus CD	75–90 minutes per session, 4 sessions total	–
Cohen et al., 2004 (United States)	Female patients with lymphoma undergoing chemotherapy; 19 were enrolled in an intervention group, and 19 were enrolled in a control group.	Tibetan yoga (mind-breathing); mixed plus CD	1 day per week; 1 self-practice session per day (no details)	7 weeks
Danhauer, Griffin, et al., 2015 (United States)	Female patients with stage I–III breast cancer undergoing chemotherapy; 22 were enrolled in an intervention group and 18 in a control group.	Integral yoga; mixed plus CD	75 minutes per week, with 45 minutes per day twice per week of self-practice	10 weeks
Dhruba et al., 2012 (United States)	Female patients with breast cancer undergoing chemotherapy; 9 were enrolled in an intervention group and 9 in a control group.	Pranayama breathing-based; mixed	60 minutes per week supervised; 20–30 minutes per day, 6 days per week, of self-practice	–
Jin et al., 2017 (China)	Female patients with breast cancer undergoing chemotherapy; 50 were enrolled in an intervention group and 50 in a control group.	Self-design; mixed plus CD	60 minutes per day, 3 days per week	16 weeks
Jong et al., 2018 (Netherlands)	Female patients with stage I–III breast cancer undergoing chemotherapy and/or radiation therapy; 47 were enrolled in an intervention group (39 completed) and 36 in a control group (29 completed).	Dru yoga; mixed plus CD	75 minutes per week; 5 minutes per day for 7 days of self-practice	12 weeks

*Continued on the next page*

**TABLE 1. Characteristics of Included Studies (N = 16) (Continued)**

Study and Location	Sample	Intervention	Duration	Endpoint
Moadel et al., 2007 (United States)	Female patients with breast cancer undergoing chemotherapy or antiestrogen therapy or radiation therapy; 108 were enrolled in an intervention group and 56 in a control group.	Hatha yoga; mixed plus CD	60 minutes per week with daily self-practice (no detail)	12 weeks
Sohl et al., 2016 (United States)	9 male and 6 female patients with colorectal cancer undergoing chemotherapy; 8 were enrolled in an intervention group (6 completed) and 7 in a control group (5 completed).	Self-design; home-base audio recording	15 minutes per day, 4 days per week	8 weeks
Taso et al., 2014 (Taiwan)	Female patients with breast cancer receiving adjuvant chemotherapy; 30 were enrolled in an intervention group and 30 in a control group.	Asanas yoga plus self-design; supervised	60 minutes per day, 2 days per week	4 weeks, 8 weeks
Vadiraja et al., 2009 (India)	Female patients with stage II–III breast cancer undergoing adjuvant radiation therapy; 44 were enrolled in an intervention group (42 completed) and 44 in a control group (33 completed).	Asanas plus pranayama yoga; mixed plus CD	60 minutes per week, 3 days per week	6 weeks
Wang et al., 2013 (China)	Female patients with breast cancer undergoing chemotherapy; 50 were enrolled in an intervention group (40 completed) and 50 in a control group (42 completed).	Self-design; mixed plus CD	50 minutes per day, 4 days per week	16 weeks
Xiang et al., 2017 (China)	Female patients with breast cancer undergoing chemotherapy; 24 were enrolled in a yoga intervention group, 21 were enrolled in a yoga plus music intervention group, and 23 were enrolled in a control group.	Self-design; mixed plus CD	40 minutes 7 times, with self-practice (no detail)	4 weeks

### Inclusion and Exclusion Criteria

The following criteria were required for study inclusion:

- Participants had to be older than age 18 years and undergoing chemotherapy and/or radiation therapy (regardless of type of drug or dose) for cancer (regardless of cancer type or stage).
- Study interventions must include yoga, regardless of the style, intensity, and duration, and be focused on yoga, including one or more sessions of posture yoga, meditation, and pranayama.
- The study must contain a routine care comparison group (e.g., standard nursing and/or psychoeducation).
- The primary outcome was CRF, measured through a standardized, validated, and reliable psychometric instrument, and the endpoint was during

or just after the conclusion of the yoga intervention. Secondary outcomes included adherence and adverse events.

- Experimental and quasiexperimental study designs, including RCTs and non-RCTs, were included.

Studies were excluded if less than 50% of the total participant population were those who received chemotherapy and/or radiation therapy and the data for which were not extracted independently; the yoga intervention was combined with other therapy, such as massage, acupuncture, drug, and nutrition; and the study had a single-arm design with no comparison group.

### Search Strategy

Eight English and Chinese databases, including PubMed®, Cochrane Library, OVID, EBSCOhost, ProQuest, CNKI, SINOMED, and Wanfang, were

searched for relevant studies from inception to July 2019. An initial limited search on PubMed and Cochrane Library was undertaken to identify articles on the topic. Medical Subject Heading (MeSH) words and text phrases were used to develop a full search strategy for the PubMed database. The search strategy, including MeSH words, text phrases, title, abstract, keywords, and index terms, were adapted for each included information source. The languages were limited to English and Chinese. The reference lists of all studies selected for critical appraisal were screened for additional relevant studies.

### Study Selection

Following the search, all identified citations were collected and uploaded into EndNote®, version 19.2, and duplicates were removed. Titles, abstracts, and the full text were assessed by two independent reviewers against the inclusion criteria. Any disagreements that arose between the reviewers at each stage of the study selection process were resolved through discussion or with a third reviewer.

### Assessment of Methodologic Quality

Eligible studies were critically appraised for methodologic quality by two independent reviewers at the study level using the standardized critical appraisal instruments from the Joanna Briggs Institute for experimental and quasiexperimental studies. The risks of bias were assessed on the following domains: random sequence generation, allocation concealment, blinding of participants, intervention and outcomes assessors, intention-to-treat (ITT) analysis, selective reporting, and attrition bias. Any disagreements were resolved through discussion or with a third reviewer.

### Data Extraction

Data were extracted by two independent reviewers using a standardized data extraction tool. The data extracted included specific details about the study characteristics (author, year, and country where the study was conducted), participant characteristics (age, gender, and type of cancer), study methods, treatment, intervention characteristics (the style of yoga and intensity [minutes per session, sessions per day, days per week, and total weeks]), and outcome characteristics (sample size, mean scores and effect size of CRF, the assessment tool, numbers of adherence, and adverse events). Authors of respective articles were contacted to request missing or additional data or for clarification where required.

### Statistical Analyses

Data from studies were pooled for statistical meta-analysis using RevMan, version 5.3. The main outcome index was CRF, which was a continuous variable. However, there were various measuring instruments. Effect sizes of CRF were expressed as weighted mean difference for the same measuring instruments, while standardized mean difference (SMD) was used to express effect sizes for different measuring instruments. The 95% confidence interval (CI) was calculated for analysis and modified as appropriate. Relevant adverse events (sprain, fall, dizziness) and adherence (defined as patient participation in 70% or more of intervention) were binary outcomes, which were summarized using odds ratio (OR).

Heterogeneity was assessed statistically using the standard chi-square and  $I^2$  tests. Subgroup analyses were performed regardless of the presence or absence of statistically significant heterogeneity for the following: (a) types of cancer; (b) types of yoga, categorized into no postures (pure meditation and/or breathing without physical exercises), postures (physical exercises without meditation or breathing), and mixed types (physical exercises with one or more of meditation and breathing); (c) intervention duration, categorized into less than eight weeks and eight weeks or greater; (d) frequency and intensity of intervention, including less than 150 minutes per week or 150 minutes or greater per week; (e) cancer type, including breast, colorectal, and lung; and (f) intervention strategies, categorized into supervised intervention, self-practicing, or mixed. Statistical analyses were performed using the random-effects model due to factors that could affect the outcomes; (e.g., different assessment instruments, yoga styles, and intervention strategies in studies) (Tufanaru et al., 2015). Sensitivity analyses were conducted to exclude studies with high heterogeneity of research methodology in subgroups, such as yoga style and variability in the ratio of random distribution. Publication bias was visually inspected using funnel plots and formally assessed using statistical tests. For all statistical analyses,  $p < 0.05$  indicated statistical significance.

## Results

### Study Selection

Seven hundred records were identified from searches of nine databases, and nine additional studies were identified from other review articles. After duplicates were removed, the titles and abstracts of 588

articles were screened against eligibility criteria and 526 were excluded, leaving 62 full-text articles. Of the 62 articles, 46 were excluded for the following reasons: post-treatment or less than 50% of the total participants received chemotherapy and/or radiation therapy (n = 13); no CRF assessment (n = 9); full text not available (n = 7); single-arm design (n = 6); articles not in English or Chinese (n = 5); related to the same process and outcome (n = 3); and no routine care comparison (n = 3). Sixteen RCTs comprised of 20 intervention groups were included in the final study (see Figure 1). No quasiexperimental studies met the inclusion criteria.

### Characteristics of Included Studies

The included 16 studies comprised a total of 1,453 patients, of whom 105 were men and 1,348 were women. The sample size of each study ranged from 15 to 164 patients. Nine studies were conducted in

the United States, three in mainland China, two in India, one in the Netherlands, and one in Taiwan. The types of cancer in the studies were breast cancer (n = 13), prostate cancer (n = 1), lymphoma (n = 1), and colorectal cancer (n = 1). Patients were receiving chemotherapy only (n = 9), radiation therapy only (n = 5), and chemotherapy and/or radiation therapy (n = 2). Interventions were conducted and compared with routine care in three studies, and one study measured CRF at two endpoints (during and after the intervention). Standard nursing was the comparator in 14 studies; in two other studies, psychoeducation was the comparator (see Table 1).

Regarding types of yoga as an intervention in the studies, 13 were mixed yoga, 2 were no posture (mind-breathing), and 2 were only posture (stretching practice). The styles of yoga were variable as well. The duration of the yoga interventions ranged from 4 to 16 weeks, the frequency of yoga

**TABLE 2. Effectiveness of Yoga Intervention on Cancer-Related Fatigue**

Study and Subgroup	Yoga			Control			Weight (%)	SMD	95% CI
	$\bar{X}$	SD	Total	$\bar{X}$	SD	Total			
Chandwani et al., 2010	1.9	0.7	30	2.5	0.8	31	6.6	-0.79	[-1.31, -0.26]
Chandwani et al., 2014 (stretching)	2.5	0.3	56	3.2	0.4	54	6.8	-1.97	[-2.43, -1.51]
Chandwani et al., 2014 (yoga)	2.9	0.3	53	3.2	0.4	54	7.1	-0.84	[-1.24, -0.45]
Chaoul et al., 2018 (stretching)	3.7	2.3	68	3.5	2.5	85	7.3	-0.84	[-0.24, -0.4]
Chaoul et al., 2018 (yoga)	3.2	2.4	74	3.5	2.5	85	7.3	-0.12	[-0.43, 0.19]
Cohen et al., 2004	3.1	1.5	19	3.1	1.5	19	6.1	0.00	[-0.64, 0.64]
Dhruva et al., 2012	5.6	2.1	9	4.8	2.5	9	4.9	0.33	[-0.6, 1.26]
Jong et al., 2018	14.6	4.5	47	14.2	4.2	36	6.9	0.09	[-0.34, 0.52]
Moadel et al., 2007	34.37	11.26	108	33.82	12.97	56	7.3	0.05	[-0.28, 0.37]
Taso et al., 2014 (4 weeks)	16.8	6.7	30	14.9	4.4	30	6.7	0.33	[-0.18, 0.84]
Taso et al., 2014 (8 weeks)	10.9	6.9	30	20.4	5	30	6.4	-1.56	[-2.14, -0.97]
Vadiraja et al., 2009	33.26	23.8	44	50.52	22.3	44	6.9	0.74	[-1.17, -0.31]
Wang et al., 2013	20.12	3.78	50	24.67	3.83	50	7	-1.19	[-1.61, -0.76]
Xiang et al., 2017 (yoga)	22.08	6.57	24	25.65	5.79	23	6.4	-0.57	[-1.15, 0.02]
Xiang et al., 2017 (yoga plus music)	21.17	5.53	22	25.65	5.79	23	6.3	-0.78	[-1.39, -0.17]
Total (95% CI) <sup>a</sup>	-	-	664	-	-	629	100	-0.52	[-0.86, -0.18]

<sup>a</sup> Heterogeneity:  $\tau^2 = 0.39$ ,  $\chi^2 = 119.61$ ,  $df = 14$  ( $p < 0.00001$ ),  $I^2 = 88\%$ ; test for overall effect:  $Z = 2.97$  ( $p = 0.003$ )  
CI—confidence interval; SMD—standard mean difference



interventions ranged from one to seven days per week, and the total duration of sessions ranged from 60 to 200 minutes per week. In most studies (n = 12), yoga was instituted as a combined supervised and self-practicing intervention; three studies conducted yoga intervention by supervision, and one study adopted a home-based strategy. The assessment tools for CRF included the Brief Fatigue Inventory (BFI) (n = 6), the Cancer Fatigue Scale (CFS) (n = 4), the Functional Assessment of Cancer Therapy–Fatigue subscale (n = 2), the Functional Assessment of Chronic Illness Therapy–Fatigue subscale (n = 1), the revised Piper Fatigue Scale (n = 1), the Multidimensional Fatigue Inventory (n = 1), and the European Organisation for the Research and

Treatment of Cancer Quality-of-Life Questionnaire–Core 30 (n = 1).

#### Quality Assessment and Risk of Bias

Randomization was used for the assignment of participants to different groups in all studies, but detailed information regarding random sequence generation was not described in three studies. Only one study performed cluster randomization, using one cluster for yoga and one for comparison. Apart from one study that performed a random assignment in a 2:1 ratio, all others were performed in a 1:1 ratio. Concealed allocation was reported in seven studies, of which two did not provide detailed information. All studies compared baseline data between the yoga

**TABLE 3. Subgroup 1: Effectiveness of Yoga on Cancer-Related Fatigue by Instrument**

Study and Subgroup	Yoga			Control			Weight (%)	SMD	95% CI
	$\bar{X}$	SD	Total	$\bar{X}$	SD	Total			
<b>Brief Fatigue Inventory</b>									
Chandwani et al., 2010	1.9	0.7	30	2.5	0.8	31	16.2	-0.6	[-0.98, -0.22]
Chandwani et al., 2014 (stretching)	2.5	0.3	56	3.2	0.4	54	17.9	-0.7	[-0.83, -0.57]
Chandwani et al., 2014 (yoga)	2.9	0.3	53	3.2	0.4	54	17.9	-0.3	[-0.43, -0.17]
Chaoul et al., 2018 (stretching)	3.7	2.3	68	3.5	2.5	85	12.1	0.2	[-0.56, 0.96]
Chaoul et al., 2018 (yoga)	3.2	2.4	74	3.5	2.5	85	12.1	-0.3	[-1.06, 0.46]
Cohen et al., 2004	3.1	1.5	19	3.1	1.5	19	10.2	0.00	[-0.95, 0.95]
Taso et al., 2014 (4 weeks)	16.8	6.7	30	14.9	4.4	30	2.2	1.9	[-0.97, 4.77]
Taso et al., 2014 (8 weeks)	10.9	6.9	30	20.4	5	30	2	-9.5	[-12.55, -6.45]
Subtotal (95% CI) <sup>a</sup>	-	-	360	-	-	388	90.5	-0.46	[-0.86, -0.06]
<b>Cancer Fatigue Scale</b>									
Wang et al., 2013	20.12	3.78	50	24.67	3.83	50	6.2	-1.19	[-1.61, -0.76]
Xiang et al., 2017 (yoga)	22.08	6.57	24	25.65	5.79	23	1.5	-3.57	[-7.11, -0.03]
Xiang et al., 2017 (yoga plus music)	21.17	5.53	22	25.65	5.79	23	1.7	-4.48	[-7.79, -1.17]
Subtotal (95% CI) <sup>b</sup>	-	-	96	-	-	96	9.5	-4.41	[-5.68, -3.14]
<b>Overall</b>									
Total (95% CI) <sup>c</sup>	-	-	664	-	-	629	100	-0.52	[-0.86, -0.18]

<sup>a</sup> Heterogeneity:  $\tau^2 = 0.18$ ,  $\chi^2 = 58.31$ ,  $df = 7$  ( $p < 0.00001$ ),  $I^2 = 88\%$ ; test for overall effect:  $Z = 2.27$  ( $p = 0.02$ )

<sup>b</sup> Heterogeneity:  $\tau^2 = 0.00$ ,  $\chi^2 = 0.25$ ,  $df = 2$  ( $p < 0.88$ ),  $I^2 = 0\%$ ; test for overall effect:  $Z = 6.82$  ( $p = 0.00001$ )

<sup>c</sup> Heterogeneity:  $\tau^2 = 0.3$ ,  $\chi^2 = 94.98$ ,  $df = 10$  ( $p < 0.00001$ ),  $I^2 = 89\%$ ; test for overall effect:  $Z = 3.64$  ( $p = 0.0003$ )

CI—confidence interval; SMD—standard mean difference

**Note.** Test for subgroup differences:  $\chi^2 = 33.98$ ,  $df = 1$  ( $p < 0.00001$ ),  $I^2 = 97.1\%$

and control groups. Of all included studies, only one reported no blinding to outcomes assessors. Two studies did not report reasons for patient attrition, and four did not describe details about the patients' adherence to the intervention. In two studies, patient attrition and reasons for attrition were significantly different between the groups. ITT analysis was performed in nine studies. None of the studies provided the registered protocol, so the risk of selective reporting was unclear. Only one study described the training of outcome assessors, whereas no studies provided the number of raters and intra-rater reliability. Pseudorandomization was performed in one study, whereas another study had assessed CRF using a measurement scale of unclear validity. These factors

might have affected the quality of trial design and resulted in biases.

#### Pooled Effectiveness of a Yoga Intervention on CRF

All 16 studies reported outcomes on CRF. However, two studies presented the data by median and quartile, one study used only graphical representation, one study did not calculate the total scores of CRF, and one study compared the difference in the value of CRF between the two groups before and after the intervention. Therefore, 11 studies involving 15 results were pooled into the meta-analysis. When comparing the total effect on CRF of yoga intervention with routine care, the pooled SMD was  $-0.52$  (95% CI  $[-0.86, -0.18]$ ), indicating that yoga was favorable

**TABLE 4. Subgroup 1: Effectiveness of Yoga on Cancer-Related Fatigue by Instrument (Sensitivity Analysis)**

Study and Subgroup	Yoga			Control			Weight (%)	SMD	95% CI
	$\bar{X}$	SD	Total	$\bar{X}$	SD	Total			
<b>Brief Fatigue Inventory</b>									
Chandwani et al., 2010	1.9	0.7	30	2.5	0.8	31	21.4	-0.6	[-0.98, -0.22]
Chandwani et al., 2014 (stretching)	2.5	0.3	56	3.2	0.4	54	-	-0.7	[-0.83, -0.57]
Chandwani et al., 2014 (yoga)	2.9	0.3	53	3.2	0.4	54	22.7	-0.3	[-0.43, -0.17]
Chaoul et al., 2018 (stretching)	3.7	2.3	68	3.5	2.5	85	-	0.2	[-0.56, 0.96]
Chaoul et al., 2018 (yoga)	3.2	2.4	74	3.5	2.5	85	17.8	-0.3	[-1.06, 0.46]
Cohen et al., 2004	3.1	1.5	19	3.1	1.5	19	15.8	0.00	[-0.95, 0.95]
Taso et al., 2014 (4 weeks)	16.8	6.7	30	14.9	4.4	30	4.5	1.9	[-0.97, 4.77]
Taso et al., 2014 (8 weeks)	10.9	6.9	30	20.4	5	30	-	-9.5	[-12.55, -6.45]
Subtotal (95% CI) <sup>a</sup>	-	-	206	-	-	219	82.3	-0.34	[-0.56, -0.13]
<b>Cancer Fatigue Scale</b>									
Wang et al., 2013	20.12	3.78	50	24.67	3.83	50	10.9	-4.55	[-6.04, -3.06]
Xiang et al., 2017 (yoga)	22.08	6.57	24	25.65	5.79	23	3.2	-3.57	[-7.11, -0.03]
Xiang et al., 2017 (yoga plus music)	21.17	5.53	22	25.65	5.79	23	3.6	-4.48	[-7.79, -1.17]
Subtotal (95% CI) <sup>b</sup>	-	-	96	-	-	96	17.7	-4.41	[-5.68, -3.14]
<b>Overall</b>									
Total (95% CI) <sup>c</sup>	-	-	302	-	-	315	100	-0.93	[-1.62, -0.25]

<sup>a</sup> Heterogeneity:  $\tau^2 = 0.01$ ,  $\chi^2 = 4.94$ ,  $df = 4$  ( $p < 0.29$ ),  $I^2 = 19\%$ ; test for overall effect:  $Z = 3.15$  ( $p = 0.002$ )

<sup>b</sup> Heterogeneity:  $\tau^2 = 0.00$ ,  $\chi^2 = 0.25$ ,  $df = 2$  ( $p < 0.88$ ),  $I^2 = 0\%$ ; test for overall effect:  $Z = 6.82$  ( $p = 0.00001$ )

<sup>c</sup> Heterogeneity:  $\tau^2 = 0.53$ ,  $\chi^2 = 44.72$ ,  $df = 7$  ( $p < 0.00001$ ),  $I^2 = 84\%$ ; test for overall effect:  $Z = 2.69$  ( $p = 0.007$ )

CI—confidence interval; SMD—standard mean difference

**Note.** Test for subgroup differences:  $\chi^2 = 38.41$ ,  $df = 1$  ( $p < 0.00001$ ),  $I^2 = 97.4\%$



and statistically significant in reducing CRF during chemotherapy/radiation therapy with a large heterogeneity ( $I^2 = 88\%$ ,  $p < 0.00001$ ). The funnel plot appeared symmetrical, suggesting a low risk of publication bias (see Table 2).

**Subgroup 1: Effectiveness on CRF with different assessment instruments:** Of the 11 studies included for meta-analysis, 6 ( $n = 748$  patients) assessed CRF using the BFI, which showed the pooled effect of  $-0.46$  (95% CI  $[-0.86, -0.06]$ ) and high heterogeneity

**TABLE 5. Subgroup 2: Effectiveness of Yoga on Cancer-Related Fatigue by Yoga Type**

Study and Subgroup	Yoga			Control			Weight (%)	SMD	95% CI
	$\bar{X}$	SD	Total	$\bar{X}$	SD	Total			
<b>Mixed</b>									
Chandwani et al., 2010	1.9	0.7	30	2.5	0.8	31	6.6	-0.79	[-1.31, -0.26]
Chandwani et al., 2014 (yoga)	2.9	0.3	53	3.2	0.4	54	7.1	-0.84	[-1.24, -0.45]
Cohen et al., 2004	3.1	1.5	19	3.1	1.5	19	6.1	0.00	[-0.64, 0.64]
Jong et al., 2018	14.6	4.5	47	14.2	4.2	36	6.9	0.09	[-0.34, 0.52]
Moadel et al., 2007	34.37	11.26	108	33.82	12.97	56	7.3	0.05	[-0.28, 0.37]
Taso et al., 2014 (4 weeks)	16.8	6.7	30	14.9	4.4	30	6.7	0.33	[-0.18, 0.84]
Taso et al., 2014 (8 weeks)	10.9	6.9	30	20.4	5	30	6.4	-1.56	[-2.14, -0.97]
Vadiraja et al., 2009	33.26	23.8	44	50.52	22.3	44	6.9	-0.74	[-1.17, -0.31]
Wang et al., 2013	20.12	3.78	50	24.67	3.83	50	7	-1.19	[-1.61, -0.76]
Xiang et al., 2017 (yoga)	22.08	6.57	24	25.65	5.79	23	6.4	-0.57	[-1.15, 0.02]
Xiang et al., 2017 (yoga plus music)	21.17	5.53	22	25.65	5.79	23	6.3	-0.78	[-1.39, -0.17]
Subtotal (95% CI) <sup>a</sup>	-	-	457	-	-	396	73.6	-0.54	[-0.89, -0.19]
<b>Mind-breathing</b>									
Chaoul et al., 2018 (yoga)	3.2	2.4	74	3.5	2.5	85	7.3	-0.12	[-0.43, 0.19]
Dhruva et al., 2012	5.6	2.1	9	4.8	2.5	9	4.9	0.33	[-0.6, 1.26]
Subtotal (95% CI) <sup>b</sup>	-	-	83	-	-	94	12.3	-0.08	[-0.37, 0.22]
<b>Posture</b>									
Chandwani et al., 2014 (stretching)	2.5	0.3	56	3.2	0.4	54	6.8	-1.97	[-2.43, -1.51]
Chaoul et al., 2018 (stretching)	3.7	2.3	68	3.5	2.5	85	7.3	0.08	[-0.24, 0.4]
Subtotal (95% CI) <sup>c</sup>	-	-	124	-	-	139	14.2	-0.94	[-2.95, 1.08]
<b>Overall</b>									
Total (95% CI) <sup>d</sup>	-	-	664	-	-	629	100	-0.52	[-0.86, -0.18]

<sup>a</sup> Heterogeneity:  $\tau^2 = 0.28$ ,  $\chi^2 = 59.33$ ,  $df = 10$  ( $p < 0.00001$ ),  $I^2 = 83\%$ ; test for overall effect:  $Z = 3.03$  ( $p = 0.002$ )

<sup>b</sup> Heterogeneity:  $\tau^2 = 0.00$ ,  $\chi^2 = 0.81$ ,  $df = 1$  ( $p < 0.37$ ),  $I^2 = 0\%$ ; test for overall effect:  $Z = 0.5$  ( $p = 0.61$ )

<sup>c</sup> Heterogeneity:  $\tau^2 = 2.07$ ,  $\chi^2 = 51.95$ ,  $df = 1$  ( $p < 0.00001$ ),  $I^2 = 98\%$ ; test for overall effect:  $Z = 0.91$  ( $p = 0.36$ )

<sup>d</sup> Heterogeneity:  $\tau^2 = 0.39$ ,  $\chi^2 = 119.61$ ,  $df = 14$  ( $p < 0.00001$ ),  $I^2 = 88\%$ ; test for overall effect:  $Z = 2.97$  ( $p = 0.003$ )

CI—confidence interval; SMD—standard mean difference

**Note.** Test for subgroup differences:  $\chi^2 = 4.35$ ,  $df = 2$  ( $p < 0.11$ ),  $I^2 = 54\%$

( $I^2 = 88\%$ ,  $p < 0.00001$ ). Two studies ( $n = 192$  patients) measured CRF using the CFS, which demonstrated the pooled effect of  $-4.41$  (95% CI  $[-5.68, 3.14]$ ) and low heterogeneity ( $I^2 = 0\%$ ,  $p = 0.88$ ) (see Table 3). The other four studies measured CRF using a different instrument; therefore, the measured data from these studies were not merged in the analysis. The difference between subgroups was statistically significant ( $I^2 = 97.1\%$ ,  $p < 0.00001$ ), indicating that the

assessment instrument was an important factor contributing to heterogeneity. A sensitivity analysis was performed for the BFI subgroup, which showed that the heterogeneity was reduced ( $I^2 = 19\%$ ,  $p = 0.29$ ) following the omission of three interventions, and the outcome appeared stable (SMD =  $-0.34$ , 95% CI  $[-0.56, -0.13]$ ) (see Table 4).

**Subgroup 2: Effectiveness on CRF by yoga type:** Of the included studies, 11 interventions ( $n = 853$

**TABLE 6. Subgroup 3: Effectiveness of Yoga on Cancer-Related Fatigue by Intervention Strategies**

Study and Subgroup	Yoga			Control			Weight (%)	SMD	95% CI
	$\bar{X}$	SD	Total	$\bar{X}$	SD	Total			
<b>Supervised</b>									
Taso et al., 2014 (4 weeks)	16.8	6.7	30	14.9	4.4	30	6.7	0.33	$[-0.18, 0.84]$
Taso et al., 2014 (8 weeks)	10.9	6.9	30	20.4	5	30	6.4	-1.56	$[-2.14, -0.97]$
Subtotal (95% CI) <sup>a</sup>	-	-	60	-	-	60	13	-0.61	$[-2.46, 1.24]$
<b>Supervised plus self-practice</b>									
Chandwani et al., 2010	1.9	0.7	30	2.5	0.8	31	6.6	-0.79	$[-1.31, -0.26]$
Chandwani et al., 2014 (stretching)	2.5	0.3	56	3.2	0.4	54	6.8	-1.97	$[-2.43, -1.51]$
Chandwani et al., 2014 (yoga)	2.9	0.3	53	3.2	0.4	54	7.1	-0.84	$[-1.24, -0.45]$
Chaoul et al., 2018 (stretching)	3.7	2.3	68	3.5	2.5	85	7.3	0.08	$[-0.24, 0.4]$
Chaoul et al., 2018 (yoga)	3.2	2.4	74	3.5	2.5	85	7.3	-0.12	$[-0.43, 0.19]$
Cohen et al., 2004	3.1	1.5	19	3.1	1.5	19	6.1	0.00	$[-0.64, 0.64]$
Dhruva et al., 2012	5.6	2.1	9	4.8	2.5	9	4.9	0.33	$[-0.6, 1.26]$
Jong et al., 2018	14.6	4.5	47	14.2	4.2	36	6.9	0.09	$[-0.34, 0.52]$
Moadel et al., 2007	34.37	11.26	108	33.82	12.97	56	7.3	0.05	$[-0.28, 0.37]$
Vadiraja et al., 2009	33.26	23.8	44	50.52	22.3	44	6.9	-0.74	$[-1.17, -0.31]$
Wang et al., 2013	20.12	3.78	50	24.67	3.83	50	7	-1.19	$[-1.61, -0.76]$
Xiang et al., 2017 (yoga)	22.08	6.57	24	25.65	5.79	23	6.4	-0.57	$[-1.15, 0.02]$
Xiang et al., 2017 (yoga plus music)	21.17	5.53	22	25.65	5.79	23	6.3	-0.78	$[-1.39, -0.17]$
Subtotal (95% CI) <sup>b</sup>	-	-	604	-	-	569	87	-0.51	$[-0.86, -0.16]$
<b>Overall</b>									
Total (95% CI) <sup>c</sup>	-	-	664	-	-	629	100	-0.52	$[-0.86, -0.18]$

<sup>a</sup> Heterogeneity:  $\tau^2 = 1.7$ ,  $\chi^2 = 22.84$ ,  $df = 1$  ( $p < 0.00001$ ),  $I^2 = 96\%$ ; test for overall effect:  $Z = 0.64$  ( $p = 0.52$ )

<sup>b</sup> Heterogeneity:  $\tau^2 = 0.35$ ,  $\chi^2 = 96.73$ ,  $df = 12$  ( $p < 0.00001$ ),  $I^2 = 88\%$ ; test for overall effect:  $Z = 2.85$  ( $p = 0.004$ )

<sup>c</sup> Heterogeneity:  $\tau^2 = 0.39$ ,  $\chi^2 = 119.61$ ,  $df = 14$  ( $p < 0.00001$ ),  $I^2 = 88\%$ ; test for overall effect:  $Z = 2.97$  ( $p = 0.003$ )

CI—confidence interval; SMD—standard mean difference

**Note.** Test for subgroup differences:  $\chi^2 = 0.01$ ,  $df = 1$  ( $p < 0.92$ ),  $I^2 = 0\%$

patients) used mixed yoga types; the pooled effect on CRF was  $-0.54$  (95% CI  $[-0.89, -0.19]$ ) with substantial heterogeneity ( $I^2 = 83\%$ ,  $p < 0.00001$ ). Two interventions implemented posture-based or no posture (mind-breathing) yoga, and the pooled effects were  $-0.94$  (95% CI  $[-2.95, 1.08]$ ,  $I^2 = 98\%$ ,  $p < 0.00001$ ) and  $-0.08$  (95% CI  $[-0.37, 0.22]$ ,  $I^2 = 0\%$ ,  $p = 0.37$ ), respectively (see Table 5).

**Subgroup 3: Effectiveness on CRF by intervention strategies:** Ten studies ( $n = 1,173$  patients) had implemented yoga with combined supervision and

self-practicing strategy. The pooled effect on CRF was  $-0.51$  (95% CI  $[-0.86, -0.16]$ ) with substantial heterogeneity ( $I^2 = 88\%$ ,  $p < 0.00001$ ) (see Table 6). Only one study instituted yoga intervention by supervised strategy.

**Subgroup 4: Effectiveness on CRF with a variable weekly duration of yoga intervention:** The weekly total durations of yoga interventions were less than 150 minutes per week in six studies ( $n = 802$  patients) and 150 minutes or greater per week in four studies ( $n = 423$  patients). The pooled effect showed that yoga

**TABLE 7. Subgroup 4: Effectiveness of Yoga on Cancer-Related Fatigue by Duration of Intervention**

Study and Subgroup	Yoga			Control			Weight (%)	SMD	95% CI
	$\bar{X}$	SD	Total	$\bar{X}$	SD	Total			
<b>Less than 150 minutes per week</b>									
Chandwani et al., 2010	1.9	0.7	30	2.5	0.8	31	7	-0.79	[-1.31, -0.26]
Chaoul et al., 2018 (stretching)	3.7	2.3	68	3.5	2.5	85	7.8	0.08	[-0.24, 0.4]
Chaoul et al., 2018 (yoga)	3.2	2.4	74	3.5	2.5	85	7.8	-0.12	[-0.43, 0.19]
Jong et al., 2018	14.6	4.5	47	14.2	4.2	36	7.4	0.09	[-0.34, 0.52]
Moadel et al., 2007	34.37	11.26	108	33.82	12.97	56	7.8	0.05	[-0.28, 0.37]
Taso et al., 2014 (4 weeks)	16.8	6.7	30	14.9	4.4	30	7.1	0.33	[-0.18, 0.84]
Taso et al., 2014 (8 weeks)	10.9	6.9	30	20.4	5	30	6.8	-1.56	[-2.14, -0.97]
Xiang et al., 2017 (yoga)	22.08	6.57	24	25.65	5.79	23	6.8	-0.57	[-1.15, 0.02]
Xiang et al., 2017 (yoga plus music)	21.17	5.53	22	25.65	5.79	23	6.7	-0.78	[-1.39, -0.17]
Subtotal (95% CI) <sup>a</sup>	-	-	433	-	-	399	65.1	-0.32	[-0.65, 0.01]
<b>150 or more minutes per week</b>									
Chandwani et al., 2014 (stretching)	2.5	0.3	56	3.2	0.4	54	7.3	-1.97	[-2.43, -1.51]
Chandwani et al., 2014 (yoga)	2.9	0.3	53	3.2	0.4	54	7.5	-0.84	[-1.24, -0.45]
Dhruva et al., 2012	5.6	2.1	9	4.8	2.5	9	5.3	0.33	[-0.6, 1.26]
Vadiraaja et al., 2009	33.26	23.8	44	50.52	22.3	44	7.4	-0.74	[-1.17, -0.31]
Wang et al., 2013	20.12	3.78	50	24.67	3.83	50	7.4	-1.19	[-1.61, -0.76]
Subtotal (95% CI) <sup>b</sup>	-	-	212	-	-	211	34.9	-0.96	[-1.52, -0.4]
<b>Overall</b>									
Total (95% CI) <sup>c</sup>	-	-	645	-	-	610	100	-0.55	[-0.91, -0.19]

<sup>a</sup> Heterogeneity:  $\tau^2 = 0.2$ ,  $\chi^2 = 41.99$ ,  $df = 8$  ( $p < 0.00001$ ),  $I^2 = 81\%$ ; test for overall effect:  $Z = 1.9$  ( $p = 0.06$ )

<sup>b</sup> Heterogeneity:  $\tau^2 = 0.34$ ,  $\chi^2 = 27.3$ ,  $df = 4$  ( $p < 0.00001$ ),  $I^2 = 85\%$ ; test for overall effect:  $Z = 3.33$  ( $p = 0.0009$ )

<sup>c</sup> Heterogeneity:  $\tau^2 = 0.4$ ,  $\chi^2 = 117.65$ ,  $df = 13$  ( $p < 0.00001$ ),  $I^2 = 89\%$ ; test for overall effect:  $Z = 3.03$  ( $p = 0.002$ )

CI—confidence interval; SMD—standard mean difference

**Note.** Test for subgroup differences:  $\chi^2 = 3.64$ ,  $df = 1$  ( $p < 0.06$ ),  $I^2 = 72.5\%$

intervention of 150 minutes or greater per week significantly reduced CRF ( $-0.96$ , 95% CI  $[-1.52, -0.4]$ ,  $I^2 = 85\%$ ,  $p < 0.0001$ ) compared to routine care. When the yoga intervention was less than 150 minutes per week, CRF was only reduced slightly and not significantly different compared to routine care ( $-0.32$ , 95% CI  $[-0.65, 0.01]$ ,  $I^2 = 81\%$ ,  $p < 0.00001$ ) (see Table 7).

**Subgroup 5: Effectiveness on CRF with different intervention durations:** The duration of intervention

in seven studies ( $n = 868$  patients) was less than eight weeks, and five studies ( $n = 425$  patients) had an intervention of eight weeks or greater. The pooled effect showed that yoga intervention of less than eight weeks reduced the CRF during chemotherapy, and the difference between the groups was statistically significant ( $-0.54$ , 95% CI  $[-0.95, -0.12]$ ,  $I^2 = 88\%$ ,  $p < 0.00001$ ). However, when comparing yoga intervention and routine care in relieving CRF for

**TABLE 8. Subgroup 5: Effectiveness of Yoga on Cancer-Related Fatigue by Total Duration of Intervention**

Study and Subgroup	Yoga			Control			Weight (%)	SMD	95% CI
	$\bar{X}$	SD	Total	$\bar{X}$	SD	Total			
<b>Less than 8 weeks</b>									
Chandwani et al., 2010	1.9	0.7	30	2.5	0.8	31	6.6	-0.79	[-1.31, -0.26]
Chandwani et al., 2014 (stretching)	2.5	0.3	56	3.2	0.4	54	6.8	-1.97	[-2.43, -1.51]
Chandwani et al., 2014 (yoga)	2.9	0.3	53	3.2	0.4	54	7.1	-0.84	[-1.24, -0.45]
Chaoul et al., 2018 (stretching)	3.7	2.3	68	3.5	2.5	85	7.3	0.08	[-0.24, 0.4]
Chaoul et al., 2018 (yoga)	3.2	2.4	74	3.5	2.5	85	7.3	-0.12	[-0.43, 0.19]
Cohen et al., 2004	3.1	1.5	19	3.1	1.5	19	6.1	0.00	[-0.64, 0.64]
Taso et al., 2014 (4 weeks)	16.8	6.7	30	14.9	4.4	30	6.7	0.33	[-0.18, 0.84]
Vadhiraja et al., 2009	33.26	23.8	44	50.52	22.3	44	6.9	-0.74	[-1.17, -0.31]
Xiang et al., 2017 (yoga)	22.08	6.57	24	25.65	5.79	23	6.4	-0.57	[-1.15, 0.02]
Xiang et al., 2017 (yoga plus music)	21.17	5.53	22	25.65	5.79	23	6.3	-0.78	[-1.39, -0.17]
Subtotal (95% CI) <sup>a</sup>	-	-	420	-	-	448	67.5	-0.54	[-0.95, -0.12]
<b>8 weeks or longer</b>									
Dhruva et al., 2012	5.6	2.1	9	4.8	2.5	9	4.9	0.33	[-0.6, 1.26]
Jong et al., 2018	14.6	4.5	47	14.2	4.2	36	6.9	0.09	[-0.34, 0.52]
Moadel et al., 2007	34.37	11.26	108	33.82	12.97	56	7.3	0.05	[-0.28, 0.37]
Taso et al., 2014 (8 weeks)	10.9	6.9	30	20.4	5	30	6.4	-1.56	[-2.14, -0.97]
Wang et al., 2013	20.12	3.78	50	24.67	3.83	50	7	-1.19	[-1.61, -0.76]
Subtotal (95% CI) <sup>b</sup>	-	-	244	-	-	181	32.5	-0.47	[-1.18, 0.23]
<b>Overall</b>									
Total (95% CI) <sup>c</sup>	-	-	664	-	-	629	100	-0.52	[-0.86, -0.18]

<sup>a</sup> Heterogeneity:  $\tau^2 = 0.39$ ,  $\chi^2 = 76.49$ ,  $df = 9$  ( $p < 0.00001$ ),  $I^2 = 88\%$ ; test for overall effect:  $Z = 2.55$  ( $p = 0.01$ )

<sup>b</sup> Heterogeneity:  $\tau^2 = 0.56$ ,  $\chi^2 = 42.81$ ,  $df = 4$  ( $p < 0.00001$ ),  $I^2 = 91\%$ ; test for overall effect:  $Z = 1.32$  ( $p = 0.19$ )

<sup>c</sup> Heterogeneity:  $\tau^2 = 0.39$ ,  $\chi^2 = 119.61$ ,  $df = 14$  ( $p < 0.00001$ ),  $I^2 = 88\%$ ; test for overall effect:  $Z = 2.97$  ( $p = 0.003$ )

CI—confidence interval; SMD—standard mean difference

**Note.** Test for subgroup differences:  $\chi^2 = 0.02$ ,  $df = 1$  ( $p < 0.88$ ),  $I^2 = 0\%$

the duration of eight weeks or greater, there was no statistically significant difference in the pooled effect ( $-0.47$ , 95% CI  $[-1.18, 0.23]$ ,  $I^2 = 91\%$ ,  $p < 0.00001$ ) (see Table 8).

**Subgroup 6: Effectiveness on CRF by cancer type:**

Ten studies examined the effectiveness of yoga on CRF in patients with breast cancer, which showed the pooled effect of  $-0.46$  (95% CI  $[-0.58, -0.35]$ ) with high heterogeneity ( $I^2 = 89\%$ ,  $p < 0.00001$ ) (see Table 9).

**Pooled Effects of Adherence to Yoga Intervention**

Fourteen studies reported the adherence rate to intervention, which ranged from 7% to 100%. One study reported 100% adherence; however, this was not estimated in the pooled result. The adherence rates of 13 studies were pooled into the meta-analysis (see Table 10), which showed a pooled OR of 0.55 (95% CI  $[0.4, 0.74]$ ) with medium heterogeneity ( $I^2 = 66\%$ ,  $p = 0.0002$ ). The funnel plot appeared asymmetrical,

**TABLE 9. Subgroup 6: Effectiveness of Yoga on Cancer-Related Fatigue by Cancer Type**

Study and Subgroup	Yoga			Control			Weight (%)	SMD	95% CI
	$\bar{X}$	SD	Total	$\bar{X}$	SD	Total			
<b>Breast cancer</b>									
Chandwani et al., 2010	1.9	0.7	30	2.5	0.8	31	4.8	-0.79	[-1.31, -0.26]
Chandwani et al., 2014 (stretching)	2.5	0.3	56	3.2	0.4	54	6.2	-1.97	[-2.43, -1.51]
Chandwani et al., 2014 (yoga)	2.9	0.3	53	3.2	0.4	54	8.3	-0.84	[-1.24, -0.45]
Chaoul et al., 2018 (stretching)	3.7	2.3	68	3.5	2.5	85	12.8	0.08	[-0.24, 0.4]
Chaoul et al., 2018 (yoga)	3.2	2.4	74	3.5	2.5	85	13.4	-0.12	[-0.43, 0.19]
Dhruva et al., 2012	5.6	2.1	9	4.8	2.5	9	1.5	0.33	[-0.6, 1.26]
Jong et al., 2018	14.6	4.5	47	14.2	4.2	36	6.9	0.09	[-0.34, 0.52]
Moadel et al., 2007	34.37	11.26	108	33.82	12.97	56	12.5	0.05	[-0.28, 0.37]
Taso et al., 2014 (4 weeks)	16.8	6.7	30	14.9	4.4	30	5	0.33	[-0.18, 0.84]
Taso et al., 2014 (8 weeks)	10.9	6.9	30	20.4	5	30	3.8	-1.56	[-2.14, -0.97]
Vadiraja et al., 2009	33.26	23.8	44	50.52	22.3	44	7	-0.74	[-1.17, -0.31]
Wang et al., 2013	20.12	3.78	50	24.67	3.83	50	7.2	-1.19	[-1.61, -0.76]
Xiang et al., 2017 (yoga)	22.08	6.57	24	25.65	5.79	23	3.8	-0.57	[-1.15, 0.02]
Xiang et al., 2017 (yoga plus music)	21.17	5.53	22	25.65	5.79	23	3.5	-0.78	[-1.39, -0.17]
Subtotal (95% CI) <sup>a</sup>	-	-	645	-	-	610	96.8	-0.46	[-0.58, -0.35]
<b>Lymphoma</b>									
Cohen et al., 2004	3.1	1.5	19	3.1	1.5	19	3.2	0.00	[-0.64, 0.64]
Subtotal (95% CI) <sup>b</sup>	-	-	19	-	-	19	3.2	0.00	[-0.64, 0.64]
<b>Overall</b>									
Total (95% CI) <sup>c</sup>	-	-	664	-	-	629	100	-0.45	[-0.56, -0.33]

<sup>a</sup> Heterogeneity:  $\chi^2 = 117.65$ ,  $df = 13$  ( $p < 0.00001$ );  $I^2 = 89\%$ ; test for overall effect:  $Z = 7.81$  ( $p = 0.00001$ )

<sup>b</sup> Heterogeneity: Not applicable; test for overall effect:  $Z = 0.00$  ( $p = 1.00$ )

<sup>c</sup> Heterogeneity:  $\chi^2 = 119.61$ ,  $df = 14$  ( $p < 0.00001$ );  $I^2 = 88\%$ ; test for overall effect:  $Z = 7.68$  ( $p = 0.00001$ )

CI—confidence interval; SMD—standard mean difference

**Note.** Test for subgroup differences:  $\chi^2 = 1.97$ ,  $df = 1$  ( $p = 0.16$ ),  $I^2 = 49.2\%$

indicating a potential risk of publication bias. Sensitivity analysis showed that there was significant heterogeneity in two studies; when removed, the pooled OR of adherence between the groups was 0.65 (95% CI [0.46, 0.94]) with low heterogeneity ( $I^2 = 8\%$ ,  $p = 0.37$ ) (see Table 11). Additional subgroup analyses were performed.

**Subgroup 1: Impact on adherence with a different intervention strategy:** Two studies involving 128 patients who had performed a yoga intervention via supervised-based strategy were examined. The pooled OR of adherence was 0.18 (95% CI [0.06, 0.52]) with low heterogeneity ( $I^2 = 0\%$ ,  $p = 0.81$ ), indicating that yoga interventions with supervision contributed to a statistically significant difference in the odds of adherence between the groups of patients undergoing chemotherapy and/or radiation therapy. Ten interventions involving 884 patients who had performed yoga by supervised-based strategy in addition to self-practicing strategy were examined. The

pooled OR of adherence was 0.8 (95% CI [0.54, 1.19]) with low heterogeneity ( $I^2 = 0\%$ ,  $p = 0.79$ ) (see Table 12).

**Subgroup 2: Impact on adherence with different weekly duration of yoga intervention:** Five interventions involving 375 patients who had performed yoga interventions 150 minutes or greater per week were examined, as were seven interventions involving 599 patients who had performed less than 150 minutes per week. One study did not report a detailed weekly duration of intervention. The pooled OR of adherence on intervention of 150 minutes or greater per week and less than 150 minutes per week was 1.09 (95% CI [0.6, 1.96]) with low heterogeneity ( $I^2 = 0\%$ ,  $p = 0.89$ ) and 0.46 (95% CI [0.28, 0.76]) with low heterogeneity ( $I^2 = 24\%$ ,  $p = 0.25$ ), respectively (see Table 13).

**Subgroup 3: Impact on adherence with different cancer types:** Ten interventions involving 891 patients with breast cancer had a pooled OR of 0.77 (95% CI

**TABLE 10. Effectiveness of Yoga Intervention on Adherence**

Study and Subgroup	Yoga		Standard Control		Weight (%)	OR	95% CI
	Events	Total	Events	Total			
Ben-Josef et al., 2017	18	35	28	33	12.2	0.19	[0.06, 0.6]
Chandwani et al., 2010	27	30	31	31	3	0.12	[0.01, 2.52]
Chandwani et al., 2014 (stretching)	52	56	48	54	3	1.63	[0.43, 6.11]
Chandwani et al., 2014 (yoga)	49	53	48	54	3.1	1.53	[0.41, 5.77]
Chaoul et al., 2018 (stretching)	59	68	79	85	8.1	0.5	[0.17, 1.48]
Chaoul et al., 2018 (yoga)	64	74	79	85	8.6	0.49	[0.17, 1.41]
Cohen et al., 2004	12	19	14	19	4.5	0.61	[0.15, 2.44]
Danhauer, Griffin, et al., 2015	12	22	10	18	4.3	0.96	[0.27, 3.36]
Dhruva et al., 2012	8	9	8	9	0.8	1	[0.05, 18.91]
Jong et al., 2018	40	47	29	36	4.2	1.38	[0.44, 4.36]
Moadel et al., 2007	32	108	44	56	35.4	1.38	[0.44, 4.36]
Sohl et al., 2016	6	8	5	7	1.2	1.2	[0.12, 11.87]
Taso et al., 2014 (8 weeks)	27	30	30	30	3	0.13	[0.01, 2.61]
Vadiraja et al., 2009	42	44	33	44	1.3	7	[1.45, 33.79]
Wang et al., 2013	40	50	42	50	7.3	0.76	[0.27, 2.12]
Total (95% CI) <sup>a</sup>	488	653	528	611	100	0.55	[0.4, 0.74]

<sup>a</sup> Heterogeneity:  $\chi^2 = 40.59$ ,  $df = 14$  ( $p = 0.0002$ ),  $I^2 = 66\%$ ; test for overall effect:  $Z = 3.93$  ( $p < 0.0001$ )  
 CI—confidence interval; OR—odds ratio



[0.51, 1.16]) with high heterogeneity ( $I^2 = 0\%$ ,  $p = 0.59$ ). Three studies involving 121 patients with prostate cancer, lymphoma, and colorectal cancer had a pooled OR of 0.36 (95% CI [0.16, 0.81]), with medium heterogeneity ( $I^2 = 29\%$ ,  $p = 0.25$ ) (see Table 14). The validity of this result was low because of the small sample size.

The results of other subgroup analyses showed no significant difference among different yoga styles and intervention durations on patient adherence. Only four of the included studies reported no adverse events associated with yoga.

## Discussion

This systematic review and meta-analysis included a reasonably adequate number of studies and total numbers of patients ( $n = 16$ , 1,453 patients), which provided evidence on the effectiveness of yoga on CRF in patients undergoing chemotherapy and/or radiation therapy. Compared with routine care, the

current analysis has demonstrated that yoga could reduce CRF, which is consistent with findings from Cramer et al. (2017), but contrary to that of Hilfiker et al. (2018) and Pan et al. (2015). Small sample sizes and high heterogeneity between included studies may have contributed to inaccuracy, inconsistency, and variability among several systematic reviews. Also, in previous reviews, additional subgroup analyses were not performed.

Subgroup analysis in the current study demonstrated that yoga type, intervention strategy, weekly and total duration, and assessment instrument were significant factors that affected the reported effectiveness of yoga on CRF. Most of the studies included in the current review instituted mixed yoga, and the current meta-analysis has confirmed the effectiveness of mixed yoga compared to posture-only (stretching) or mind-breathing yoga. Patients with cancer experience significant fatigue, of which the root cause is often

**TABLE 11. Effectiveness of Yoga Intervention on Adherence (Total, Sensitivity Analysis)**

Study and Subgroup	Yoga		Standard Control		Weight (%)	OR	95% CI
	Events	Total	Events	Total			
Ben-Josef et al., 2017	18	35	28	33	19.2	0.19	[0.06, 0.6]
Chandwani et al., 2010	27	30	31	31	4.8	0.12	[0.01, 2.52]
Chandwani et al., 2014 (stretching)	52	56	48	54	4.8	1.63	[0.43, 6.11]
Chandwani et al., 2014 (yoga)	49	53	48	54	4.9	1.53	[0.41, 5.77]
Chaoul et al., 2018 (stretching)	59	68	79	85	12.7	0.5	[0.17, 1.48]
Chaoul et al., 2018 (yoga)	64	74	79	85	13.6	0.49	[0.17, 1.41]
Cohen et al., 2004	12	19	14	19	7.1	0.61	[0.15, 2.44]
Danhauer, Griffin, et al., 2015	12	22	10	18	6.9	0.96	[0.27, 3.36]
Dhruva et al., 2012	8	9	8	9	1.2	1	[0.05, 18.91]
Jong et al., 2018	40	47	29	36	6.7	1.38	[0.44, 4.36]
Moadel et al., 2007	32	108	44	56	-	1.38	[0.44, 4.36]
Sohl et al., 2016	6	8	5	7	1.8	1.2	[0.12, 11.87]
Taso et al., 2014 (8 weeks)	27	30	30	30	4.7	0.13	[0.01, 2.61]
Vadiraja et al., 2009	42	44	33	44	-	7	[1.45, 33.79]
Wang et al., 2013	40	50	42	50	11.5	0.76	[0.27, 2.12]
Total (95% CI) <sup>a</sup>	414	501	451	511	100	0.65	[0.46, 0.94]

<sup>a</sup> Heterogeneity:  $\chi^2 = 13.04$ ,  $df = 12$  ( $p = 0.37$ ),  $I^2 = 8\%$ ; test for overall effect:  $Z = 2.29$  ( $p < 0.02$ )  
CI—confidence interval; OR—odds ratio

multidimensional, including physical, emotional, and/or cognitive factors. Mixed yoga could improve the physical function and mental state of patients undergoing cancer treatment.

Contrary to other reviews (Lipsett et al., 2017; Meneses-Echávez et al., 2015; Velthuis et al., 2010), the current meta-analysis revealed that supervision in addition to self-practicing strategy was more effective than supervision alone. Such inconsistency of findings may

be related to the fact that the majority of the patients in the included studies were outpatients or part-time inpatients, who not only have been experiencing treatment-induced fatigue but also other life events, which may make them prefer to have flexible arrangements of yoga schedule and location of intervention. However, supervision of treatment at regular intervals would appear to be more conducive to providing physical and mental health support, which would improve patient

**TABLE 12. Subgroup 1: Effectiveness of Yoga Intervention on Adherence**

Study and Subgroup	Yoga		Standard Control		Weight (%)	OR	95% CI
	Events	Total	Events	Total			
<b>Supervised</b>							
Ben-Josef et al., 2017	18	35	28	33	19.2	0.19	[0.06, 0.6]
Taso et al., 2014 (8 weeks)	27	30	30	30	4.7	0.13	[0.01, 2.61]
Subtotal (95% CI) <sup>a</sup>	45	65	58	63	23.9	0.18	[0.06, 0.52]
<b>Supervised plus self-practice</b>							
Chandwani et al., 2010	27	30	31	31	4.8	0.12	[0.01, 2.52]
Chandwani et al., 2014 (stretching)	52	56	48	54	4.8	1.63	[0.43, 6.11]
Chandwani et al., 2014 (yoga)	49	53	48	54	4.9	1.53	[0.41, 5.77]
Chaoul et al., 2018 (stretching)	59	68	79	85	12.7	0.5	[0.17, 1.48]
Chaoul et al., 2018 (yoga)	64	74	79	85	13.6	0.49	[0.17, 1.41]
Cohen et al., 2004	12	19	14	19	7.1	0.61	[0.15, 2.44]
Danhauer, Griffin, et al., 2015	12	22	10	18	6.9	0.96	[0.27, 3.36]
Dhruva et al., 2012	8	9	8	9	1.2	1	[0.05, 18.91]
Jong et al., 2018	40	47	29	36	6.7	1.38	[0.44, 4.36]
Moadel et al., 2007	32	108	44	56	-	NE	NE
Sohl et al., 2016	6	8	5	7	1.8	1.2	[0.12, 11.87]
Vadiraja et al., 2009	42	44	33	44	-	NE	NE
Wang et al., 2013	40	50	42	50	11.5	0.76	[0.27, 2.12]
Subtotal (95% CI) <sup>b</sup>	369	436	393	448	76.1	0.8	[0.54, 1.19]
<b>Overall</b>							
Total (95% CI) <sup>c</sup>	414	501	451	511	100	0.65	[0.46, 0.94]

<sup>a</sup> Heterogeneity:  $\chi^2 = 0.06$ ,  $df = 1$  ( $p = 0.81$ ),  $I^2 = 0\%$ ; test for overall effect:  $Z = 3.13$  ( $p < 0.002$ )

<sup>b</sup> Heterogeneity:  $\chi^2 = 6.29$ ,  $df = 10$  ( $p = 0.79$ ),  $I^2 = 0\%$ ; test for overall effect:  $Z = 1.08$  ( $p < 0.28$ )

<sup>c</sup> Heterogeneity:  $\chi^2 = 13.04$ ,  $df = 12$  ( $p = 0.37$ ),  $I^2 = 8\%$ ; test for overall effect:  $Z = 2.29$  ( $p < 0.02$ )

CI—confidence interval; NE—not estimable; OR—odds ratio

**Note.** Test for subgroup differences:  $\chi^2 = 6.63$ ,  $df = 1$  ( $p = 0.01$ ),  $I^2 = 84.9\%$

adherence with treatment. Again, a small sample size in the subgroup of supervised strategy may contribute to the inconsistency of the results between meta-analyses.

Several previous reviews have concluded that a high volume of exercise does not relieve CRF in patients undergoing treatment for cancer (Carayol et al., 2015; Kessels et al., 2018; Schwartz et al., 2001; Velthuis et al., 2010), although the definition of high-volume exercise has been inconsistent among the studies. In the current analysis, yoga with a weekly duration of 150 minutes or greater per week

and a total intervention duration of less than eight weeks appeared to be the most effective in reducing CRF.

In the reviewed studies, the adopted assessment instrument consisted of six scales, in which the reliability and validity have been verified in patients with cancer (Gebremariam et al., 2018; Paramita et al., 2016). The current authors did not merge the data from four of the six scales in one study. The BFI was the most commonly used assessment tool for CRF in the included studies. The current meta-analysis has

**TABLE 13. Subgroup 2: Effectiveness of Yoga Intervention on Adherence**

Study and Subgroup	Yoga		Standard Control		Weight (%)	OR	95% CI
	Events	Total	Events	Total			
<b>Less than 150 minutes</b>							
Ben-Josef et al., 2017	18	35	28	33	20.7	0.19	[0.06, 0.6]
Chandwani et al., 2010	27	30	31	31	5.2	0.12	[0.01, 2.52]
Chaoul et al., 2018 (stretching)	59	68	79	85	13.7	0.5	[0.17, 1.48]
Chaoul et al., 2018 (yoga)	64	74	79	85	14.7	0.49	[0.17, 1.41]
Jong et al., 2018	40	47	29	36	7.2	1.38	[0.44, 4.36]
Moadel et al., 2007	32	108	44	56	-	NE	NE
Sohl et al., 2016	6	8	5	7	2	1.2	[0.12, 11.87]
Taso et al., 2014 (8 weeks)	27	30	30	30	5.1	0.13	[0.01, 2.61]
Subtotal (95% CI) <sup>a</sup>	241	292	281	307	68.5	0.46	[0.28, 0.76]
<b>150 minutes or more</b>							
Chandwani et al., 2014 (stretching)	52	56	48	54	5.2	1.63	[0.43, 6.11]
Chandwani et al., 2014 (yoga)	49	53	48	54	5.3	1.53	[0.41, 5.77]
Danhauer, Griffin, et al., 2015	12	22	10	18	7.4	0.96	[0.27, 3.36]
Dhruva et al., 2012	8	9	8	9	1.3	1	[0.05, 18.91]
Vadiraja et al., 2009	42	44	33	44	-	NE	NE
Wang et al., 2013	40	50	42	50	12.4	0.76	[0.27, 2.12]
Subtotal (95% CI) <sup>b</sup>	161	190	156	185	31.5	1.09	[0.6, 1.96]
<b>Overall</b>							
Total (95% CI) <sup>c</sup>	402	482	437	492	100	0.66	[0.45, 0.96]
<sup>a</sup> Heterogeneity: $\chi^2 = 7.87$ , $df = 6$ ( $p = 0.25$ ), $I^2 = 24\%$ ; test for overall effect: $Z = 3.03$ ( $p = 0.002$ )							
<sup>b</sup> Heterogeneity: $\chi^2 = 1.11$ , $df = 4$ ( $p = 0.89$ ), $I^2 = 0\%$ ; test for overall effect: $Z = 0.28$ ( $p < 0.78$ )							
<sup>c</sup> Heterogeneity: $\chi^2 = 13.02$ , $df = 11$ ( $p = 0.29$ ), $I^2 = 15\%$ ; test for overall effect: $Z = 2.19$ ( $p = 0.03$ )							
CI—confidence interval; NE—not estimable; OR—odds ratio							
<b>Note.</b> Test for subgroup differences: $\chi^2 = 4.77$ , $df = 1$ ( $p = 0.03$ ), $I^2 = 79.1\%$							

shown that the effectiveness of yoga intervention in the subgroup analyzed with the BFI appeared lower but more precise than interventions analyzed using the CFS.

The heterogeneity was substantial in the current meta-analysis, which could not be reduced by subgroup analysis. However, in the BFI subgroup, three dubious interventions have been identified in the sensitivity analyses; when these were omitted, the

heterogeneity was reduced and the outcome became steady. Causes of heterogeneity in the current analysis would likely be multifactorial, including different assessment instruments, yoga type (only stretching intervention in two studies) (Chandwani et al., 2014; Chaoul et al., 2018), and the outcome assessment on CRF following yoga intervention being too early (four weeks) (Taso et al., 2014), which were the major sources of heterogeneity.

**TABLE 14. Subgroup 3: Effectiveness of Yoga Intervention on Adherence**

Study and Subgroup	Yoga		Standard Control		Weight (%)	OR	95% CI
	Events	Total	Events	Total			
<b>Breast cancer</b>							
Chandwani et al., 2010	27	30	31	31	4.8	0.12	[0.01, 2.52]
Chandwani et al., 2014 (stretching)	52	56	48	54	4.8	1.63	[0.43, 6.11]
Chandwani et al., 2014 (yoga)	49	53	48	54	4.9	1.53	[0.41, 5.77]
Chaoul et al., 2018 (stretching)	59	68	79	85	12.7	0.5	[0.17, 1.48]
Chaoul et al., 2018 (yoga)	64	74	79	85	13.6	0.49	[0.17, 1.41]
Danhauer et al., 2015	12	22	10	18	6.9	0.96	[0.27, 3.36]
Dhruva et al., 2012	8	9	8	9	1.2	1	[0.05, 18.91]
Jong et al., 2018	40	47	29	36	6.7	1.38	[0.44, 4.36]
Moadel et al., 2007	32	108	44	56	-	NE	NE
Taso et al., 2014 (8 weeks)	27	30	30	30	4.7	0.13	[0.01, 2.61]
Vadiraja et al., 2009	42	44	33	44	-	NE	NE
Wang et al., 2013	40	50	42	50	11.5	0.76	[0.27, 2.12]
Subtotal (95% CI) <sup>a</sup>	378	439	404	452	71.9	0.77	[0.51, 1.16]
<b>Non-breast cancer</b>							
Ben-Josef et al., 2017	18	35	28	33	19.2	0.19	[0.06, 0.6]
Cohen et al., 2004	12	19	14	19	7.1	0.61	[0.15, 2.44]
Sohl et al., 2016	6	8	5	7	1.8	1.2	[0.12, 11.87]
Subtotal (95% CI) <sup>b</sup>	36	62	47	59	28.1	0.36	[0.16, 0.81]
<b>Overall</b>							
Total (95% CI) <sup>c</sup>	414	501	451	511	100	0.65	[0.46, 0.94]
<sup>a</sup> Heterogeneity: $\chi^2 = 7.49$ , $df = 9$ ( $p = 0.59$ ), $I^2 = 0\%$ ; test for overall effect: $Z = 1.25$ ( $p = 0.21$ ) <sup>b</sup> Heterogeneity: $\chi^2 = 2.81$ , $df = 2$ ( $p = 0.25$ ), $I^2 = 29\%$ ; test for overall effect: $Z = 2.49$ ( $p = 0.01$ ) <sup>c</sup> Heterogeneity: $\chi^2 = 13.04$ , $df = 12$ ( $p = 0.37$ ), $I^2 = 8\%$ ; test for overall effect: $Z = 2.29$ ( $p = 0.02$ ) CI—confidence interval; NE—not estimable; OR—odds ratio <b>Note.</b> Test for subgroup differences: $\chi^2 = 2.71$ , $df = 1$ ( $p = 0.1$ ), $I^2 = 63.1\%$							

### Patient Adherence With Yoga Intervention

Patient adherence is one of the most important factors when determining the effectiveness of an intervention. To the authors' knowledge, few reviews have performed a narrative-only analysis of patient adherence to an exercise intervention (Buffart et al., 2012; Meneses-Echávez et al., 2015; Velthuis et al., 2010). The current study represents the first meta-analysis examining patient adherence to yoga interventions while undergoing cancer treatment. Factors for patient attrition included schedule conflicts, movement restrictions, transportation issues, lack of time, difficulty in attending regularly, changing treatment regimens, and other health-related problems. In the current study, several factors may have affected patient adherence. Of note, the adherence of patients involved in a supervised-based yoga intervention less than 150 minutes per week was significantly lower, in parallel with the effectiveness on CRF. This suggests that a supervised-based format in addition to self-practicing strategy might offer more flexibility and improve adherence to outpatient visits. An optimal intervention duration might correspond to improved CRF, leading to increased persistence and confidence in the practice of yoga.

When patient adherence was analyzed in the current study, significant heterogeneity was noted among three studies; one study performed random assignment in a 2:1 ratio and the others performed in a 1:1 random assignment. Patient adherence might be increased when the yoga intervention is instituted in an inpatient setting, as was the case in the study by Vadiraja et al. (2009).

### Quality of Evidence

The overall risk of bias of the included studies was medium to low. Of the 16 studies, 12 provided detailed information regarding random sequence generation, of which allocation concealment and detailed information were reported in five studies (Chakrabarty et al., 2015; Cohen et al., 2004; Moadel et al., 2007; Taso et al., 2014; Vadiraja et al., 2009). This resulted in a moderate risk of selection bias. Blinding is an important principle in the RCT setting but is difficult to implement when the investigators need to know the intervention and when CRF is a patient-reported outcome. Therefore, none of the studies implemented blinding. Although downgrading of the evidence did not occur as a result of this, the risk of performance and detection bias remained notable. Nine studies were classified as low risk of attrition bias because ITT analysis was performed and adequate

---

### KNOWLEDGE TRANSLATION

- As a low-intensity exercise, yoga appears beneficial in alleviating cancer-related fatigue in patients undergoing chemotherapy and/or radiation therapy.
  - To increase patient adherence with this intervention, a mixed yoga with supervised practice in addition to a self-practicing strategy is required.
  - Compared to posture-only (stretching) or mind-breathing yoga, mixed yoga is more effective for relieving fatigue in patients undergoing chemotherapy and/or radiation therapy.
- 

information regarding reasons for patient attrition were provided (Ben-Josef et al., 2017; Chakrabarty et al., 2015; Chandwani et al., 2010, 2014; Chaoul et al., 2018; Danhauer, Griffin, et al., 2015; Sohl et al., 2016; Taso et al., 2014). None of the studies provided the registered protocol; therefore, the risk of selective reporting remains unclear.

### Limitations

The current study had several limitations. First, there were very small sample sizes (less than 30 participants) in four of the included studies, which could increase the risk of bias. Second, there was high heterogeneity in the analyses of pooled effects of yoga on CRF, which could not be eliminated or reduced by performing subgroup analysis. Third, various instruments were used in the studies for the assessment of CRF. Therefore, performing a pooled analysis of data became difficult. Fourth, detailed information about participants who completed the yoga intervention was reported in only nine studies (Chandwani et al., 2010, 2014; Chaoul et al., 2018; Cohen et al., 2004; Danhauer, Griffin, et al., 2015; Jong et al., 2018; Moadel et al., 2007; Taso et al., 2014; Vadiraja et al., 2009), and data on attrition had to be extracted to replace the actual participation of other studies, which could influence the ultimate results.

### Implications for Practice and Research

Despite the methodologic limitations, the practice of yoga, particularly mixed yoga that combines with posture and either mind and/or breathing control exercise, can be recommended as an additional strategy in the management of CRF in patients undergoing chemotherapy and/or radiation therapy. Supervised yoga practice in addition to self-practicing strategy using yoga manuals or DVDs might be beneficial in increasing patient adherence and improving the

effectiveness of yoga intervention. As a low-intensity exercise, yoga practice of 150 minutes or greater per week is safe and effective. Additional research consisting of adequate sample sizes, variety of cancer types other than breast cancer, a clear objective of assessment and outcome, inclusivity of blinding (at least the outcome assessors), and assessment of patient adherence to yoga intervention is required. Future studies should focus on strengthening the supervision of yoga exercise as management for CRF in the outpatient setting.

## Conclusion

This study provides valuable evidence on the effect of yoga interventions on CRF in patients undergoing chemotherapy and/or radiation therapy. Yoga is a safe and effective tool in this regard. Mixed yoga and supervised in addition to self-practicing strategy might be more beneficial in increasing patient adherence and relieving CRF. Additional rigorous studies are required to explore an optimal strategy for yoga intervention.

---

**Jihong Song, MA**, is a PhD student in the School of Nursing at Xi'an Jiaotong University Health Science Center in Xi'an, Shaanxi, and a teacher in the School of Nursing at Fujian Medical University in FuZhou, Fujian; **Tao Wang, MA**, is a PhD student in the School of Nursing at Xi'an Jiaotong University Health Science Center and a teacher at Hainan Medical University in Haikou, Hainan; **Yujie Wang, MA**, is a PhD student in the School of Nursing at Peking University in Beijing; **Rong Li, MA**, is a nurse in Qilu Hospital of Shandong University in JiNan, Shandong; **Sitian Niu, BSN**, and **Litao Zhuo, BSN**, are students in the School of Nursing at Fujian Medical University; and **Qian Guo, MA**, is a research assistant, and **Xiaomei Li, PhD**, is a professor, both in the School of Nursing at Xi'an Jiaotong University Health Science Center, all in China. X. Li can be reached at roselee8825@163.com, with copy to ONFEditor@ons.org. (Submitted July 2020. Accepted October 10, 2020.)

This study was supported by the Natural Science Foundation of Fujian Province, China (grant no. 2017J01815).

X. Li contributed to the conceptualization and design. T. Wang, Y. Wang, R. Li, Niu, Zhuo, and Guo completed the data collection. Song, T. Wang, Y. Wang, and X. Li provided analysis. Song, Guo, and X. Li contributed to the manuscript preparation.

## REFERENCES

American Association for Cancer Research. (2018). *AACR cancer progress report 2018*. [https://cancerprogressreport.aacr.org/wp-content/uploads/sites/2/2020/09/AACR\\_CPR\\_2018.pdf](https://cancerprogressreport.aacr.org/wp-content/uploads/sites/2/2020/09/AACR_CPR_2018.pdf)

- Ben-Josef, A.M., Chen, J., Wileyto, P., Doucette, A., Bekelman, J., Christodouleas, J., . . . Vapiwala, N. (2017). Effect of Eischens yoga during radiation therapy on prostate cancer patient symptoms and quality of life: A randomized phase II trial. *International Journal of Radiation Oncology, Biology, Physics*, 98(5), 1036–1044. <https://doi.org/10.1016/j.ijrobp.2017.03.043>
- Buffart, L.M., van Uffelen, J.G.Z., Riphagen, I.I., Brug, J., van Mechelen, W., Brown, W.J., & Chinapaw, M.J.M. (2012). Physical and psychosocial benefits of yoga in cancer patients and survivors, a systematic review and meta-analysis of randomized controlled trials. *BMC Cancer*, 12, 559. <https://doi.org/10.1186/1471-2407-12-559>
- Carayol, M., Bernard, P., Boiché, J., Riou, F., Mercier, B., Cousson-Gélie, F., . . . Ninot, G. (2013). Psychological effect of exercise in women with breast cancer receiving adjuvant therapy: What is the optimal dose needed? *Annals of Oncology*, 24(2), 291–300. <https://doi.org/10.1093/annonc/mds342>
- Carayol, M., Delpierre, C., Bernard, P., & Ninot, G. (2015). Population-, intervention- and methodology-related characteristics of clinical trials impact exercise efficacy during adjuvant therapy for breast cancer: A meta-regression analysis. *Psycho-Oncology*, 24(7), 737–747. <https://doi.org/10.1002/pon.3727>
- Chakrabarty, J., Vidyasagar, M., Fernandes, D., Joisa, G., Varghese, P., & Mayya, S. (2015). Effectiveness of pranayama on cancer-related fatigue in breast cancer patients undergoing radiation therapy: A randomized controlled trial. *International Journal of Yoga*, 8(1), 47–53. <https://doi.org/10.4103/0973-6131.146062>
- Chandwani, K.D., Perkins, G., Nagendra, H.R., Raghuram, N.V., Spelman, A., Nagarathna, R., . . . Cohen, L. (2014). Randomized, controlled trial of yoga in women with breast cancer undergoing radiotherapy. *Journal of Clinical Oncology*, 32(10), 1058–1065. <https://doi.org/10.1200/JCO.2012.48.2752>
- Chandwani, K.D., Thornton, B., Perkins, G.H., Arun, B., Raghuram, N.V., Nagendra, H.R., . . . Cohen, L. (2010). Yoga improves quality of life and benefit finding in women undergoing radiotherapy for breast cancer. *Journal of the Society for Integrative Oncology*, 8(2), 43–55. <https://doi.org/10.2310/7200.2010.0002>
- Chaoul, A., Milbury, K., Spelman, A., Basen-Engquist, K., Hall, M.H., Wei, Q., . . . Cohen, L. (2018). Randomized trial of Tibetan yoga in patients with breast cancer undergoing chemotherapy. *Cancer*, 124(1), 36–45. <https://doi.org/10.1002/cncr.30938>
- Cohen, L., Warneke, C., Fouladi, R.T., Rodriguez, M.A., & Chaoul-Reich, A. (2004). Psychological adjustment and sleep quality in a randomized trial of the effects of a Tibetan yoga intervention in patients with lymphoma. *Cancer*, 100(10), 2253–2260. <https://doi.org/10.1002/cncr.20236>
- Cramer, H., Lauche, R., Klose, P., Lange, S., Langhorst, J., & Dobos, G.J. (2017). Yoga for improving health-related quality of life, mental health and cancer-related symptoms in women diagnosed with breast cancer. *Cochrane Database of Systematic*



- Reviews, 1, CD010802. <https://doi.org/10.1002/14651858.CD010802.pub2>
- Cramer, H., Pokhrel, B., Fester, C., Meier, B., Gass, F., Lauche, R., . . . Langhorst, J. (2016). A randomized controlled bicenter trial of yoga for patients with colorectal cancer. *Psycho-Oncology*, 25(4), 412–420. <https://doi.org/10.1002/pon.3927>
- Danhauer, S.C., Addington, E.L., Cohen, L., Sohl, S.J., Van Puymbroeck, M., Albinati, N.K., & Culos-Reed, S.N. (2019). Yoga for symptom management in oncology: A review of the evidence base and future directions for research. *Cancer*, 125(12), 1979–1989. <https://doi.org/10.1002/cncr.31979>
- Danhauer, S.C., Addington, E.L., Sohl, S.J., Chaoul, A., & Cohen, L. (2015). Evidence supports incorporating yoga alongside conventional cancer treatment for women with breast cancer. *Breast Diseases*, 26(3), 189–193. <https://doi.org/10.1016/j.breastdis.2015.07.036>
- Danhauer, S.C., Griffin, L.P., Avis, N.E., Sohl, S.J., Jesse, M.T., Addington, E.L., . . . Shaw, E. (2015). Feasibility of implementing a community-based randomized trial of yoga for women undergoing chemotherapy for breast cancer. *Journal of Community and Supportive Oncology*, 13(4), 139–147. <https://doi.org/10.12788/jcso.0125>
- Dhruva, A., Miaskowski, C., Abrams, D., Acree, M., Cooper, B., Goodman, S., & Hecht, F.M. (2012). Yoga breathing for cancer chemotherapy-associated symptoms and quality of life: Results of a pilot randomized controlled trial. *Journal of Alternative and Complementary Medicine*, 18(5), 473–479. <https://doi.org/10.1089/acm.2011.0555>
- Felbel, S., Meerpohl, J.J., Monsef, I., Engert, A., & Skoetz, N. (2014). Yoga in addition to standard care for patients with haematological malignancies. *Cochrane Database of Systematic Reviews*, 6, CD010146. <https://doi.org/10.1002/14651858.CD010146.pub2>
- Gebremariam, G.T., Anshabo, A.T., Tigeneh, W., & Engidawork, E. (2018). Validation of the Amharic version of the Brief Fatigue Inventory for Assessment of Cancer-Related Fatigue in Ethiopian cancer patients. *Journal of Pain and Symptom Management*, 56(2), 264–272. <https://doi.org/10.1016/j.jpainsymman.2018.04.015>
- Harder, H., Parlour, L., & Jenkins, V. (2012). Randomised controlled trials of yoga interventions for women with breast cancer: A systematic literature review. *Supportive Care in Cancer*, 20(12), 3055–3064. <https://doi.org/10.1007/s00520-012-1611-8>
- Hilfiker, R., Meichtry, A., Eicher, M., Nilsson Balfe, L., Knols R.H., Verra M.L., & Taeymans J. (2018). Exercise and other non-pharmaceutical interventions for cancer-related fatigue in patients during or after cancer treatment: A systematic review incorporating an indirect-comparisons meta-analysis. *British Journal of Sports Medicine*, 52(10), 651–658. <https://doi.org/10.1136/bjsports-2016-096422>
- Jin, C., Wang, L., & Wang, B. (2017). Effects of yoga on cancer-related fatigue and quality of life in breast cancer patients with chemotherapy. *Chinese Journal of Integrative Nursing*, 3(4), 12–15. <https://doi.org/10.11997/nitcwm.201704004>
- Jong, M.C., Boers, I., Schouten van der Velden, A.P., van der Meij, S., Göker, E., Timmer-Bonte, A.N.J.H., & van Wietmarschen, H.A. (2018). A randomized study of yoga for fatigue and quality of life in women with breast cancer undergoing (neo) adjuvant chemotherapy. *Journal of Alternative and Complementary Medicine*, 24(9–10), 942–953. <https://doi.org/10.1089/acm.2018.0191>
- Kessels, E., Husson, O., & van der Feltz-Cornelis, C.M. (2018). The effect of exercise on cancer-related fatigue in cancer survivors: A systematic review and meta-analysis. *Neuropsychiatric Disease and Treatment*, 14, 479–494. <https://doi.org/10.2147/NDT.S150464>
- Lipsett, A., Barrett, S., Haruna, F., Mustian, K., & O'Donovan, A. (2017). The impact of exercise during adjuvant radiotherapy for breast cancer on fatigue and quality of life: A systematic review and meta-analysis. *Breast*, 32, 144–155. <https://doi.org/10.1016/j.breast.2017.02.002>
- Lötzke, D., Wiedemann, F., Rodrigues Recchia, D., Ostermann, T., Sattler, D., Ettl, J., . . . Büsing, A. (2016). Iyengar-Yoga compared to exercise as a therapeutic intervention during (neo)adjuvant therapy in women with stage I–III breast cancer: Health-related quality of life, mindfulness, spirituality, life satisfaction, and cancer-related fatigue. *Evidence-Based Complementary and Alternative Medicine*, 2016, 5931816. <https://doi.org/10.1155/2016/5931816>
- Meneses-Echávez, J.F., González-Jiménez, E., & Ramirez-Velez, R. (2015). Effects of supervised multimodal exercise interventions on cancer-related fatigue: Systematic review and meta-analysis of randomized controlled trials. *BioMed Research International*, 2015, 328636. <https://doi.org/10.1155/2015/328636>
- Moadel, A.B., Shah, C., Wylie-Rosett, J., Harris, M.S., Patel, S.R., Hall, C.B., & Sparano, J.A. (2007). Randomized controlled trial of yoga among a multiethnic sample of breast cancer patients: Effects on quality of life. *Journal of Clinical Oncology*, 25(28), 4387–4395. <https://doi.org/10.1200/JCO.2006.06.6027>
- Moher, D., Shamseer, L., Clarke, M., Ghersi, D., Liberati, A., Petticrew, M., . . . Stewart, L.A. (2015). Preferred Reporting Items for Systematic Reviews and Meta-Analysis protocols (PRISMA-P) 2015 statement. *Systematic Reviews*, 4, 1. <https://doi.org/10.1186/2046-4053-4-1>
- National Comprehensive Cancer Network. (2018). NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines®): Cancer-related fatigue [v.2.2018]. [https://www.nccn.org/professionals/physician\\_gls/default.aspx](https://www.nccn.org/professionals/physician_gls/default.aspx)
- Pan, Y., Yang, K., Wang, Y., Zhang, L., & Liang, H. (2015). Could yoga practice improve treatment-related side effects and quality of life for women with breast cancer? A systematic review and meta-analysis. *Asia-Pacific Journal of Clinical Oncology*, 13(2), e79–e95. <https://doi.org/10.1111/ajco.12329>
- Paramita, N., Nusdwinuringtyas, N., Nuhonni, S.A., Atmakusuma, T.D., Ismail, R.I., Mendoza, T.R., & Cleeland, C.S. (2016). Validity and reliability of the Indonesian version of the Brief Fatigue

- Inventory in cancer patients. *Journal of Pain and Symptom Management*, 52(5), 744–751.
- Sadja, J., & Mills, P.J. (2013). Effects of yoga interventions on fatigue in cancer patients and survivors: A systematic review of randomized controlled trials. *Explore*, 9(4), 232–243. <https://doi.org/10.1016/j.explore.2013.04.005>
- Schmitz, K.H., Courneya, K.S., Matthews, C., Demark-Wahnefried, W., Galvão, D.A., Pinto, B.M., . . . Schwartz, A.L. (2010). American College of Sports Medicine roundtable on exercise guidelines for cancer survivors. *Medicine and Science in Sports and Exercise*, 42(7), 1409–1426. <https://doi.org/10.1249/MSS.0b013e3181e0c112>
- Schwartz, A.L., Mori, M., Gao, R., Nail, L.M., & King, M.E. (2001). Exercise reduces daily fatigue in women with breast cancer receiving chemotherapy. *Medicine and Science in Sports and Exercise*, 33(5), 718–723.
- Smith, K.B., & Pukall, C.F. (2009). An evidence-based review of yoga as a complementary intervention for patients with cancer. *Psycho-Oncology*, 18(5), 465–475.
- Sohl, S.J., Danhauer, S.C., Birdee, G.S., Nicklas, B.J., Yacoub, G., Aklilu, M., & Avis, N.E. (2016). A brief yoga intervention implemented during chemotherapy: A randomized controlled pilot study. *Complementary Therapies in Medicine*, 25, 139–142. <https://doi.org/10.1016/j.ctim.2016.02.003>
- Streeter, C.C., Gerbarg, P.L., Saper, R.B., Ciraulo, D.A., & Brown, R.P. (2012). Effects of yoga on the autonomic nervous system, gamma-aminobutyric-acid, and allostasis in epilepsy, depression, and post-traumatic stress disorder. *Medical Hypotheses*, 78(5), 571–579. <https://doi.org/10.1016/j.mehy.2012.01.021>
- Taso, C.J., Lin, H.S., Lin, W.L., Chen, S.M., Huang, W.T., & Chen, S.W. (2014). The effect of yoga exercise on improving depression, anxiety, and fatigue in women with breast cancer: A randomized controlled trial. *Journal of Nursing Research*, 22(3), 155–164. <https://doi.org/10.1097/jnr.000000000000044>
- Tian, L., Lu, H.J., Lin, L., & Hu, Y. (2016). Effects of aerobic exercise on cancer-related fatigue: A meta-analysis of randomized controlled trials. *Supportive Care in Cancer*, 24(2), 969–983. <https://doi.org/10.1007/s00520-015-2953-9>
- Tolia, M., Tsoukalas, N., Nikolaou, M., Mosa, E., Nazos, I., Poultsidi, A., . . . Kyrgias, G. (2018). Utilizing yoga in oncologic patients treated with radiotherapy: Review. *Indian Journal of Palliative Care*, 24(3), 355–358.
- Tomlinson, D., Diorio, C., Beyene, J., & Sung, L. (2014). Effect of exercise on cancer-related fatigue: A meta-analysis. *American Journal of Physical Medicine and Rehabilitation*, 93(8), 675–686. <https://doi.org/10.1097/phm.000000000000083>
- Tufanaru, C., Munn, Z., Stephenson, M., & Aromataris, E. (2015). Fixed or random effects meta-analysis? Common methodological issues in systematic reviews of effectiveness. *International Journal of Evidence-Based Healthcare*, 13(3), 196–207.
- Vadiraja, H.S., Rao, M.R., Nagarathna, R., Nagendra, H.R., Rekha, M., Vanitha, N., . . . Rao, N. (2009). Effects of yoga program on quality of life and affect in early breast cancer patients undergoing adjuvant radiotherapy: A randomized controlled trial. *Complementary Therapies in Medicine*, 17(5–6), 274–280. <https://doi.org/10.1016/j.ctim.2009.06.004>
- Velthuis, M.J., Agasi-Idenburg, S.C., Aufdemkampe, G., & Wittink, H.M. (2010). The effect of physical exercise on cancer-related fatigue during cancer treatment: A meta-analysis of randomised controlled trials. *Clinical Oncology*, 22(3), 208–221. <https://doi.org/10.1016/j.clon.2009.12.005>
- Wang, G., Wang, S., Jiang, P., & Zeng, C. (2013). The effect of yoga on cancer related fatigue in breast cancer patients with chemotherapy. *Journal of Central South University(Medical Sciences)*, 39(10), 1077–1082. <https://doi.org/10.11817/j.issn.1672-7347.2014.10.016>
- World Health Organization. (2018). Cancer. Key facts. <https://www.who.int/en/news-room/fact-sheets/detail/cancer>
- Wu, W., Yang, F., Yang, S., Ye, C., Li, W., Li, X., & Dai, Q. (2017). Longitudinal study on the correlation between symptoms and quality of life among nasopharyngeal carcinoma patients undergoing radiotherapy. *Chinese Journal of Modern Nursing*, 23(16), 2100–2104. <https://doi.org/10.3760/cma.j.issn.1674-2907.2017.16.006>
- Xiang, D., Wang, M., Wang, H., Liu, J., & Liu, J. (2017). Intervention effect of yoga combined with music relaxation training on cancer related fatigue in breast cancer patients undergoing chemotherapy. *Chinese Journal of Modern Nursing*, 23(2), 184–187. <https://doi.org/10.3760/cma.j.issn.1674-2907.2017.02.009>
- Zuo, X.L., Li, Q., Gao, F., Yang, L., & Meng, F.J. (2016). Effects of yoga on negative emotions in patients with breast cancer: A meta-analysis of randomized controlled trials. *International Journal of Nursing Sciences*, 3(3), 299–306. <https://doi.org/10.1016/j.ijnss.2016.07.009>