

Frailty in Patients With Hematologic Malignancies and Those Undergoing Transplantation: A Scoping Review

Lathika Mohanraj, PhD, RN, BMTCN®, Lana Sargent, PhD, RN, FNP-C, GNP-BC, Roy Brown, MLIS, AHIP, and Theresa Swift-Scanlan, PhD, RN, FAAN

PROBLEM IDENTIFICATION: Although frailty is an important parameter in treatment planning and in predicting prognosis and overall survival among patients with hematologic malignancies and recipients of hematopoietic stem cell transplantation, frailty assessment tools are not standardized in clinical care settings. The purpose of this article is to provide an overview of the literature on frailty assessment tools in these patient populations.

LITERATURE SEARCH: A systematic search of CINAHL®, Embase®, MEDLINE®, PubMed®, and Web of Science was performed using keywords and controlled vocabulary for the concepts “bone marrow transplants,” “hematologic neoplasms,” and “frailty.”

DATA EVALUATION: Extracted data included study type, diagnosis, transplantation status, frailty tools used, and outcome measures.

SYNTHESIS: A systematic search resulted in 24 studies that met the inclusion criteria. There were significant differences in how various groups define and assess frailty.

IMPLICATIONS FOR PRACTICE: Addressing the lack of standardized frailty assessments will assist healthcare providers to routinely integrate frailty measures in clinical assessments to identify those at risk for poor outcomes, improving personalized care.

KEYWORDS hematopoietic stem cell transplantation; frailty; comorbidity; hematologic cancers

ONF, 48(3), 291-307.

DOI 10.1188/21.ONF.291-307

Frailty is considered “a medical syndrome with multiple causes and contributors that is characterized by diminished strength and endurance, and reduced physiologic function that increases an individual’s vulnerability for developing increased dependency and/or death” (Morley et al., 2013, p. 392). The progression from being healthy to frail is often silent and typically manifests in the clinical setting as an unexpected decline in an individual’s physical and cognitive state (Searle & Rockwood, 2015; Song et al., 2014); however, there is no clear consensus on how to operationalize the measurement of physical frailty as a vulnerable state (Pel-Littel et al., 2009). Frailty is a common clinical syndrome in older adults that increases the risk for poor health outcomes, including hospitalization, falls, and increased mortality (Perna et al., 2017). Although common in aging populations, the condition of frailty is increasingly being recognized as having a significant impact on clinical outcomes in multiple diseases and disorders such as cancer, diabetes, and heart failure (El Assar et al., 2019; Walston et al., 2018). As such, a number of specific tools have been developed to measure frailty, either as a separate construct or as one of several comorbidities in the context of a particular health state. For example, in the field of hematologic cancers and hematopoietic stem cell transplantation (HSCT), frailty has been defined, studied, and assessed in various ways. The purpose of this scoping review is to chart the state of the science for frailty assessment and describe frailty screening tools used in patients with hematologic cancers, with a focus on recipients of HSCT.


Hematologic malignancies are cancers that are initiated in the blood-forming cells of the body, such as the bone marrow and cells of the immune

system. Some examples of blood cancers are leukemias, lymphomas, and multiple myeloma. HSCT is a process by which abnormal blood-forming stem cells are replaced by healthy cells, potentially serving as a curative treatment for hematologic cancers, myelodysplastic syndrome, and other blood disorders. Type of malignancy, disease severity, and donor availability determine whether an individual receives an autologous or an allogeneic transplantation. The patient's own cells are used in an autologous transplantation, whereas in allogeneic HSCT, cells are harvested from a donor. The disease process and other factors (e.g., patient age, existing comorbidities) determine the conditioning chemotherapy regimen that patients receive prior to transplantation (Phelan et al., 2020). Treatment planning, goals, and decisions regarding aggressive chemotherapy regimens for treating hematologic cancers include eligibility determinants such as chronologic age, comorbidities, and physical performance status. To proceed to HSCT, individuals

with hematologic malignancy are expected to be in remission. Regardless of the transplantation process, these individuals have already been exposed to intense therapies to treat their cancer, with some receiving multiple regimens, that unfortunately increase their risk for frailty. The sequelae from distinct disease processes with unique clinical courses, combined with the potential for treatment-related toxicity, emphasize the need for additional assessments in patients with hematologic malignancies as well as patients undergoing HSCT (Abel & Klepin, 2018).

In addition, advancements in treatment modalities and reduced-intensity chemotherapy regimens have led to an increase in the number of patients eligible for HSCT. Despite increases in the number of novel agents and treatment regimens, there has been a scarcity of specific guidance for treatment decisions that integrate characteristics of the malignancy, as well as the patient's individual characteristics, prior therapies, and frailty. This is particularly relevant among the older adult population, where the majority of blood cancers occur and frailty is more prevalent (Abel & Klepin, 2018). The rise in the number of patients eligible for HSCT requires healthcare providers to update existing assessments to include new, relevant criteria, such as frailty, as part of developing a comprehensive risk-stratification strategy. Pidala et al. (2013) found that 92% of providers predominantly consider age as a dominant driver for predicting HSCT eligibility. In cancers such as diffuse large B-cell lymphoma, myelodysplastic syndrome, and multiple myeloma, it has been suggested that older adult patients be assessed using a multifaceted approach that involves measures of comorbidities, functional and cognitive capabilities, emotional health, nutritional status, and physiologic age (based on functional and biologic factors) for personalized therapy and predictions of therapy tolerance (Abel & Klepin, 2018; Morrison et al., 2015). Although chronologic age is an easier and important criterion to assess, it is critical to understand and acknowledge that existing comorbidities, functional and cognitive decline, and other physiologic measures need to be included in HSCT preassessments. These multidimensional parameters can also aid in monitoring prognosis, predicting therapy tolerance, and providing more personalized therapy.

In individuals with cancer, frailty may indicate chemotherapy intolerance and an increased risk of postoperative complications (Handforth et al., 2015; Rosko et al., 2015). Identification of frail patients



will help providers modify the treatment regimen so that it is suitable for these patients and intervene in a timely manner (e.g., with physical therapy) that could improve physical functioning and quality of life (Wiskemann et al., 2011). This scoping review

systematically maps literature that has used a frailty assessment or screening tool in the context of hematologic cancers and HSCT. The goal is to compile and examine data that will serve as a precursor for identifying the most promising tool(s), both in clinical

TABLE 1. Characteristics of Studies Involving Patients Undergoing HSCT (N = 11)

Study	Design and Sample	Diagnosis and Transplantation Status
Offidani et al., 2012	<ul style="list-style-type: none"> ■ Retrospective analysis ■ N = 266, median age of 73 years (range = 38-90 years) 	<ul style="list-style-type: none"> ■ MM ■ 30% autologous transplantation
Muffly et al., 2013	<ul style="list-style-type: none"> ■ Prospective pilot study ■ N = 166, median age of 58 years (range = 50-73 years) 	<ul style="list-style-type: none"> ■ AML, MDS, NHL, CML, CLL, ALL, and others ■ Allogeneic transplantation
Holmes et al., 2014	<ul style="list-style-type: none"> ■ Prospective study ■ N = 50, mean age of 65.4 years (range = 60-73 years) 	<ul style="list-style-type: none"> ■ AML, MDS, CLL, and others ■ Allogeneic transplantation
Muffly et al., 2014	<ul style="list-style-type: none"> ■ Prospective study ■ N = 203, mean age of 59 years 	<ul style="list-style-type: none"> ■ AML, MDS, NHL, CML, CLL, ALL, and others ■ Allogeneic transplantation
Arora et al., 2016	<ul style="list-style-type: none"> ■ Cohort study ■ N = 998, mean age of 42.5 years 	<ul style="list-style-type: none"> ■ Severe aplastic anemia, CML, AML, HL, NHL, ALL, MM, and others ■ 44% autologous transplantation; 46% allogeneic sibling donor; 10% matched unrelated donor
Buckstein et al., 2016	<ul style="list-style-type: none"> ■ Prospective cohort study ■ N = 445, median age of 71 years (range = 65-79 years) 	<ul style="list-style-type: none"> ■ MDS, CMML ■ 3.6% allogeneic transplantation
Milani et al., 2016	<ul style="list-style-type: none"> ■ Retrospective and prospective data ■ N = 351, median age of 65 years 	<ul style="list-style-type: none"> ■ Newly diagnosed MM ■ 39% autologous transplantation
Eissa et al., 2017	<ul style="list-style-type: none"> ■ Secondary data analysis of a prospective, longitudinal cohort study ■ N = 112, mean age of 28.4 years (SD = 5.9) 	<ul style="list-style-type: none"> ■ Hematologic malignancies ■ 29% autologous transplantation; 71% allogeneic transplantation
Engelhardt et al., 2017	<ul style="list-style-type: none"> ■ Training set and validation set ■ N = 801, median age of 63 years (28% aged 66-75 years, 13% aged older than 75 years) 	<ul style="list-style-type: none"> ■ MM ■ 57% autologous transplantation
Rosko et al., 2019	<ul style="list-style-type: none"> ■ Prospective cohort study ■ N = 100, median age of 59.5 years (range = 36-75 years) 	<ul style="list-style-type: none"> ■ MM ■ Autologous transplantation
Sakatoku et al., 2019	<ul style="list-style-type: none"> ■ Retrospective analysis ■ N = 118, median age of 73 years (range = 65-78 years) 	<ul style="list-style-type: none"> ■ MDS ■ Allogeneic transplantation

ALL—acute lymphoblastic leukemia; AML—acute myeloid leukemia; CLL—chronic lymphocytic leukemia; CML—chronic myeloid leukemia; CMML—chronic myelomonocytic leukemia; HL—Hodgkin lymphoma; HSCT—hematopoietic stem cell transplantation; MDS—myelodysplastic syndrome; MM—multiple myeloma; NHL—non-Hodgkin lymphoma

trials and in real-world clinical practice. These data will serve as a foundation to standardize assessment and guide implementation to eventually improve clinical outcomes.

Methods

Electronic Search

For this scoping review, a systematic search for published English-language literature was conducted along

TABLE 2. Characteristics of Studies Involving Patients With Hematologic Malignancy (N = 13)

Study	Design and Sample	Diagnosis
Corsetti et al., 2011	<ul style="list-style-type: none"> ■ Cohort study ■ N = 31, median age of 72 years 	AML
Marchesi et al., 2013	<ul style="list-style-type: none"> ■ Retrospective study ■ N = 73, median age of 78 years (range = 75–78 years) 	B-cell NHL
Budziszewska et al., 2015	<ul style="list-style-type: none"> ■ Prospective study ■ N = 509; median age of 68 years in fit group (range = 61–87 years), median age of 74 years in frail group (range = 61–93 years) 	AML
Okuyama et al., 2015	<ul style="list-style-type: none"> ■ Cross-sectional, observational study ■ 33 of 106 patients with MM, median age of 74 years 	MM, lymphomas
Palumbo et al., 2015	<ul style="list-style-type: none"> ■ Combined analysis of 3 prospective trials ■ N = 869; median age of 74 years, 46% of patients aged older than 75 years 	Newly diagnosed MM ineligible for transplantation
Park et al., 2015	<ul style="list-style-type: none"> ■ Prospective, observational study ■ N = 70, median age of 73.5 years 	Newly diagnosed aggressive NHL
Engelhardt et al., 2016	<ul style="list-style-type: none"> ■ Prospective validation study ■ N = 125; median age of 63 years, 26% aged 66–74 years, 15% aged older than 75 years 	MM
Velghe et al., 2016	<ul style="list-style-type: none"> ■ Prospective cohort study ■ N = 59, mean age of 77.3 years (SD = 4.8) 	AML, NHL, MDS, MM
Wilson et al., 2016	<ul style="list-style-type: none"> ■ Lifetime cohort study ■ N = 862, mean age of 31.3 years (range = 18.4–59.7 years) 	ALL
Ferrat et al., 2017	<ul style="list-style-type: none"> ■ Prospective cohort study ■ N = 763, 8% with hematologic malignancies; mean age of 80 years 	Solid tumors, hematologic malignancies
Zhong et al., 2017	<ul style="list-style-type: none"> ■ Retrospective study ■ N = 628, median age of 58 years 	Newly diagnosed MM
Hshieh et al., 2018	<ul style="list-style-type: none"> ■ Prospective, observational cohort study ■ Frailty assessment alone, n = 360; both cognitive screening tests, n = 341 of 360; mean age of 79.8 years (range = 75–79 years) 	Leukemia, myeloma, lymphoma
Molica et al., 2019	<ul style="list-style-type: none"> ■ Prospective cohort study ■ N = 108, median age of 71 years (range = 65–90 years) 	CLL

ALL—acute lymphoblastic leukemia; AML—acute myeloid leukemia; CLL—chronic lymphocytic leukemia; MDS—myelodysplastic syndrome; MM—multiple myeloma; NHL—non-Hodgkin lymphoma

with a search of the reference lists of the retrieved publications to identify published articles addressing hematologic malignancies, HSCT, and frailty. The databases searched from inception through November 19, 2018, were CINAHL[®], Embase[®], MEDLINE[®], PubMed[®], and Web of Science. The search used a combination of keywords and controlled vocabulary for the concepts “bone marrow transplants,” “hematologic neoplasms,” and “frailty” that were adapted to each database. In addition, a hand search of reference lists of articles was used to discover publications not identified in the database searches. A total of 711 articles were found, with 565 left to review after deduplication. Title and abstract screening of articles resulted in the exclusion of 380 articles. The remaining 185 full articles were assessed for eligibility. During the study selection process for this scoping review, the current authors identified cohort studies that addressed frailty in hematologic malignancies and stated “HSCT” in the article abstract or body and, therefore, met the inclusion criteria for this review. A total of 24 articles was included in this scoping review. Of these, 11 articles involved recipients of HSCT, and 13 articles included studies in individuals with hematologic malignancies. The search results and study selection process are illustrated in Figure 1, which depicts the PRISMA (Preferred Reporting Items for Systematic Reviews and Meta-Analyses) flow diagram for this scoping review (Tricco et al., 2018).

Study Selection

All references were screened, and duplicates were removed. Two of the current authors (L.M., L.S.) conducted the title and abstract review, as well as the full-text review. Any disagreements between the reviewers were resolved by consensus between the reviewers. Articles that did not involve any frailty or geriatric assessments, tools, or questionnaires were excluded. Review articles, Delphi studies, editorial opinions, expert articles, and pharmaceutical intervention studies were excluded during the selection process.

Data Collection and Analysis

The following information was extracted using a standardized data collection form: author, year of publication, diagnosis, sample population, transplantation type, and frailty assessment tool. The results were summarized in both text and tabular forms. Results were grouped in a tabular manner based on the study population: (a) patients who received an HSCT and (b) patients with hematologic malignancies. Tables 1 and 2 show characteristics of studies

involving patients undergoing HSCT and with hematologic malignancies, respectively. Tables 3 and 4 detail frailty assessment tools used in HSCT and hematologic malignancy, respectively.

Results

Tools for Frailty Screening

The use of frailty as a marker to better understand and stratify patients slated to receive HSCT and at-risk patients with blood cancers is slowly gathering momentum. However, given the differences in how frailty is defined, measured, and applied to treatment decision-making and eligibility in this population, there is an urgent need to identify standardized, reliable tools, as well as to implement measures in a consistent manner. Frailty assessment tools that have been used in the reviewed literature among patients undergoing HSCT and with hematologic cancer are summarized in Table 5. Twenty-four studies met the established criteria.

Frailty in Patients Undergoing HSCT

The current authors set out to determine how frailty involving patients undergoing HSCT was defined in the literature and the tools that were used to assess frailty in these studies. Eleven studies assessed frailty in patients undergoing HSCT, and almost half of these studies were in patients with a median age of older than 60 years. The majority of the studies used a prospective study design, followed by retrospective analysis. Frailty was assessed either by a single independent tool or by a combination of measures. Although eight different tools were used to assess frailty in these 11 studies, highlighting the heterogeneity in frailty assessments in patients undergoing HSCT, these studies indicated that frailty (as defined by each study) is associated with poor prognosis, increased mortality, decreased progression-free survival, or poor performance status. Some of the frailty tools used were Fried’s frailty phenotype (Cardiovascular Health Study [CHS] frailty index), the Clinical Frailty Scale (CFS), the Vulnerable Elders Survey-13 (VES-13), vulnerability score, the revised Myeloma Comorbidity Index (R-MCI), the International Myeloma Working Group (IMWG) score, G8, and geriatric assessments (Engelhardt et al., 2017; Holmes et al., 2014; Offidani et al., 2012; Rosko et al., 2019; Sakatoku et al., 2019).

Cardiovascular Health Study Frailty Index

The most commonly used CHS frailty index has two “frailty phenotypic” categories described as

TABLE 3. Frailty Assessment Tools in HSCT (N = 11)

Study	Frailty Tool and Components	Other Tools	Outcomes Measures
Offidani et al., 2012	Vulnerability score <ul style="list-style-type: none"> Age, performance status, renal insufficiency, bone fracture, cytopenia, comorbidity Combinatorial 	<ul style="list-style-type: none"> Within frailty assessment: World Health Organization Performance Status, CCI 	Vulnerability score distinguishes fit from unfit patients, regardless of age.
Muffly et al., 2013	CHS frailty index <ul style="list-style-type: none"> Independent 	<ul style="list-style-type: none"> Within frailty assessment: HGS Within study: CGA: ADLs, CCI, CIRS-G, HCT-CI, IADLs, Kaplan-Feinstein Scale, SF-36[®] MCS, SF-36 PCS, Zubrod PS 	CHS frailty index incorporated as part of CGA; increased frailty connected with worse performance status; recommend implementing CGA
Holmes et al., 2014	VES-13, G8, CHS frailty index <ul style="list-style-type: none"> Independent/combinatorial 	<ul style="list-style-type: none"> Within frailty assessment: G8 (HCT-CI, MNA, SPPB, ADLs, IADLs, MOS, HADS-A, HADS-D, clock draw test, trail making test); VES-13 (self-reported health, ADLs, IADLs) 	G8: higher sensitivity for abnormal GA; VES-13: higher specificity for abnormal GA; G8: correlated with CHS frailty index
Muffly et al., 2014	CHS frailty index <ul style="list-style-type: none"> Independent 	<ul style="list-style-type: none"> Within frailty assessment: HGS Within study: GA domains in addition to frailty: ADLs, CIRS-G, HCT-CI, IADLs, SF-36 MCS, SF-36 PCS 	CHS frailty index incorporated into CGA; no significant univariate associations between frailty and 2-year OS, nonrelapse mortality, and relapse; combining IADLs with HCT-CI may serve as a simple stratification tool.
Arora et al., 2016	CHS frailty index <ul style="list-style-type: none"> Independent 	-	Frailty was associated with increased risk of subsequent mortality after adjusting for relevant prognostic factors.
Buckstein et al., 2016	CFS (9-point) <ul style="list-style-type: none"> Independent 	<ul style="list-style-type: none"> Within study: SPPB, CCI, MDS-CI, IADLs 	When frailty and comorbidity are added on to IPSS, score improved by 30% and 5%, respectively.
Milani et al., 2016	IMWG frailty score, NT-proBNP <ul style="list-style-type: none"> Age, performance status, comorbidity, NT-proBNP Independent 	<ul style="list-style-type: none"> Within study: ADLs, IADLs, CCI 	Multivariate analysis of age, performance status, and NT-proBNP were independent predictors of survival.
Eissa et al., 2017	CHS frailty index <ul style="list-style-type: none"> Independent 	<ul style="list-style-type: none"> Within frailty assessment: SF-36, HGS 	Frailty was associated with an increase in risk for severe, disabling, or life-threatening pulmonary disease.
Engelhardt et al., 2017	R-MCI <ul style="list-style-type: none"> Age, impaired organ function, cytogenetics, performance status, frailty (measured by CHS frailty index) 	<ul style="list-style-type: none"> Within frailty assessment: KFI, CHS frailty index 	Frailty was incorporated as part of the R-MCI; median OS and PFS decreased from fit to frail patients.
Rosko et al., 2019	GA (11 assessments) <ul style="list-style-type: none"> Weight, physical function (physician report, self-report, objective), psychosocial metrics Combinatorial 	<ul style="list-style-type: none"> Within frailty assessment: KPS, HAP MAS, HAP AAS, BFI, HGS, SPPB, HADS-A, HADS-D, MOS-SSS, 3MS Within study: HCT-CI 	Univariate associations were observed with individual components of GA. Use of tools can aid the transplantation decision-making process and prevent morbidity.

Continued on the next page

TABLE 3. Frailty Assessment Tools in HSCT (N = 11) (Continued)

Study	Frailty Tool and Components	Other Tools	Outcomes Measures
Sakatoku et al., 2019	CFS (9-point) ■ Independent	■ Within study: CCI, MDS-CI	Recommendation to combine frailty and comorbidity with IPSS-R for precise prediction of OS

ADLs—activities of daily living; BFI—Brief Fatigue Inventory; CCI—Charlson Comorbidity Index; CFS—Clinical Frailty Scale; CGA—Comprehensive Geriatric Assessment; CHS—Cardiovascular Health Study; CIRSG—Cumulative Illness Rating Scale—Geriatric; GA—geriatric assessment; HADS-A—Hospital Anxiety and Depression Scale—Anxiety; HADS-D—Hospital Anxiety and Depression Scale—Depression; HAP AAS—Human Activity Profile Adjusted Activity Score; HAP MAS—Human Activity Profile Maximum Activity Score; HCT-CI—Hematopoietic Cell Transplantation—Specific Comorbidity Index; HGS—hand grip strength; HSCT—hematopoietic stem cell transplantation; IADLs—instrumental activities of daily living; IMWG—International Myeloma Working Group; IPSS—International Prognostic Scoring System; IPSS-R—Revised International Prognostic Scoring System; KFI—Kaplan–Feinstein Index; KPS—Karnofsky Performance Status; MCS—mental component summary; MDS-CI—Myelodysplastic Syndrome—Specific Comorbidity Index; MNA—Mini Nutritional Assessment; MOS—Medical Outcomes Study; MOS-SSS—Medical Outcomes Study—Social Support Survey; NT-proBNP—N-terminal-pro hormone B-type natriuretic peptide; OS—overall survival; PCS—physical component summary; PFS—progression-free survival; R-MCI—revised Myeloma Comorbidity Index; SPPB—Short Physical Performance Battery; 3MS—Modified Mini Mental State; VES-13—Vulnerable Elders Survey-13; Zubrod PS—Zubrod Performance Status

Note. Independent indicates that the tool is used as a known single tool for frailty; combinatorial indicates that the tool is used as a combination of tools, with frailty assessed as a part of that.

the pre-frail and frail states that are each defined by the severity of the symptom cluster consisting of fatigue, decreased physical activity, weight loss, decreased grip strength, and slowed gait. This index has concurrent and predictive validity for outcomes that include incident disease, hospitalization, falls, disability, and mortality in community-dwelling older adults (Fried et al., 2001). Although this scale has been widely used, there is heterogeneity in how specific components in this scale are measured. For instance, weight loss is measured as either unintentional weight loss in the past year or low lean muscle mass. Scores on the CHS frailty index in non-older adult patients undergoing HSCT (aged 18–64 years) approached those seen among older adults and showed a four-fold higher prevalence than those of their siblings, also suggesting that frailty in patients undergoing HSCT can occur across the lifespan, irrespective of age (Arora et al., 2016). Increased frailty has been associated with worse performance status; increased risk of severe, disabling, or life-threatening pulmonary disease; and increased risk of mortality (Arora et al., 2016; Eissa et al., 2017; Muffly et al., 2014). This tool has been used independently or incorporated as a part of geriatric assessments and the R-MCI (the R-MCI includes age, frailty, and cytogenetics) (Domm et al., 2014).

Comprehensive Geriatric Assessment

In 2005, an International Society of Geriatric Oncology taskforce recommended use of the Comprehensive Geriatric Assessment (CGA) for older adults with cancer to improve functional status and survival. Aging

in older adults is associated with significant decline in physical performance, leading to disability, loss of independence, and adverse clinical outcomes. The CGA assesses multiple domains, such as functional and mental health status, comorbidities, social support, and environmental and functional limitations; each domain is evaluated by various tools. However, frailty as a construct is not specifically assessed as a domain within the CGA. Although it is considered the gold standard for identifying older adults at high risk for adverse outcomes, the CGA has not been extensively evaluated or routinely used for patients undergoing transplantation (Extermann & Hurria, 2007; Muffly et al., 2014). This could be because of the perception that patients undergoing HSCT are typically younger than the traditional older adult population or the notion that only “fit” older adults undergo transplantation.

Nevertheless, a prospective pilot study of the CGA administered prior to allogeneic HSCT was conducted, with the intent of determining the prevalence and severity of comorbidities, vulnerabilities, and feasibility of the CGA in transplantation-eligible older adults. In the pilot study, use of the CHS frailty index score as a component of the CGA showed associations with performance status. The study recommended use of the CGA in older recipients of HSCT prior to transplantation, but also suggested functional and comorbidity assessments as simple risk stratification tools (Muffly et al., 2013, 2014).

Geriatric Assessments

Multiple studies have considered the use of select domains of the CGA to assess frailty and refer to them

TABLE 4. Frailty Assessment Tools in Hematologic Malignancy (N = 13)

Study	Frailty Tool and Components	Other Tools	Outcomes Measures
Corsetti et al., 2011	CGA (3 assessments) ■ Functional status, comorbidity ■ Combinatorial	■ Within frailty assessment: CIRS, ADLs, IADLs	Frailty and/or impairments in IADLs did not significantly affect survival in patients with AML.
Marchesi et al., 2013	CGA (4 assessments) ■ Age, functional status, comorbidity, presence or absence of geriatric syndrome (dementia, delirium, depression, incontinence, falls, osteoporosis, neglect and abuse, failure to thrive) ■ Combinatorial	■ Within frailty assessment: CIRS-G, ADLs	Patients classified as frail by the CGA showed poor OS.
Budziszewska et al., 2015	ECOG-PS, CCI ■ Performance status, comorbidity ■ Combinatorial	-	Stratifying based on ECOG-PS and CCI together increased the number of older adult patients who could be treated with the intention of inducing remission.
Okuyama et al., 2015	VES-13 ■ Combinatorial	■ Within frailty assessment: age, self-reported health, ADLs, IADLs ■ Within study: CIRS-G, PHQ-9 (anhedonia is a component of this), MMSE (delayed recall is a component of this)	The addition of anhedonia to VES-13-negative patients improves sensitivity and negative predictive value of frailty.
Palumbo et al., 2015	GA (3 assessments) ■ Functional status, comorbidity ■ Combinatorial	■ Within frailty assessment: ADLs, IADLs, CCI	Predicts mortality and risk of toxicity in older adult patients with melanoma
Park et al., 2015	GFI ■ Independent	■ Within study: MNA-SF, Korean version of MMSE, Korean geriatric depression scale, GFI	Poor GFI associated with chemotherapy discontinuation within 12 weeks. MNA-SF predicts tolerability to multi-agent chemotherapy and OS in older adult patients with aggressive NHL.
Engelhardt et al., 2016	GA (6 assessments) ■ Functional status, comorbidity ■ Combinatorial	■ Within frailty assessment: ADLs, IADLs, CCI, HCT-CI, KFI, R-MCI ■ Within study: IMWG	Because CCI is part of IMWG, it is recommended to use IMWG and R-MCI. R-MCI and IMWG have a strong prognostic value for functional decline and OS.
Velghe et al., 2016	CGA, G8, HGS ■ CGA: performance status, depression, cognitive impairment, nutrition, falls ■ Independent/combinatorial	■ Within frailty assessment: ADLs, IADLs, GDS-4, MMSE, MNA-SF	HGS was significantly higher in patients with a normal CGA compared to those with an abnormal CGA. HGS was significantly associated with abnormal G8 in women but not in men.
Wilson et al., 2016	CHS frailty index ■ Low muscle mass, exhaustion, low energy expenditure, slowness, muscle weakness ■ Independent	■ Within frailty assessment: vitality subscale of MOS, SF-36®, HGS	Frailty was associated with growth hormone deficiency-mediated decrease in bone mass density.

Continued on the next page

TABLE 4. Frailty Assessment Tools in Hematologic Malignancy (N = 13) (Continued)

Study	Frailty Tool and Components	Other Tools	Outcomes Measures
Ferrat et al., 2017	Balducci, SIOG 1, SIOG 2, LC typology <ul style="list-style-type: none"> ■ Combinations of age, functional status, comorbidity, depression, cognitive impairment ■ Combinatorial 	<ul style="list-style-type: none"> ■ Within frailty assessment: ADLs, IADLs, CIRS-G, ECOG-PS, MMSE 	Discrimination was best with SIOG 1 for 1-year mortality. SIOG 1 had the best classification agreement with Balducci and LC typology.
Zhong et al., 2017	GA (3 assessments) <ul style="list-style-type: none"> ■ Age, functional status, comorbidity ■ Combinatorial 	<ul style="list-style-type: none"> ■ Within frailty assessment: ADLs, IADLs, CCI ■ Within study: ISS 	Frail patients with ISS-II or ISS-III had the worst OS.
Hshieh et al., 2018	CHS frailty index, CFS <ul style="list-style-type: none"> ■ Independent 	<ul style="list-style-type: none"> ■ Within study: CIB, delayed recall, CCI 	Frailty by CHS frailty index correlated with executive dysfunction and/or impaired working memory.
Molica et al., 2019	GA (3 assessments) <ul style="list-style-type: none"> ■ Functional status, comorbidity ■ Combinatorial 	<ul style="list-style-type: none"> ■ Within frailty assessment: ADLs, IADLs, CIRS 	Frail and intermediate-fit patients are at a higher risk of dying than fit patients with chronic lymphocytic leukemia.

ADLs—activities of daily living; AML—acute myeloid leukemia; CCI—Charlson Comorbidity Index; CFS—Clinical Frailty Scale; CGA—Comprehensive Geriatric Assessment; CHS—Cardiovascular Health Study; CIB—clock in the box; CIRS—Cumulative Illness Rating Scale; CIRS-G—Cumulative Illness Rating Scale—Geriatric; ECOG-PS—Eastern Cooperative Oncology Group Performance Status; GA—geriatric assessment; GDS-4—4-item Geriatric Depression Scale; GFI—Groningen Frailty Index; HCT-CI—Hematopoietic Cell Transplantation—Specific Comorbidity Index; HGS—hand grip strength; IADLs—instrumental activities of daily living; IMWG—International Myeloma Working Group; ISS—International Staging System; KFI—Kaplan–Feinstein Index; MMSE—Mini-Mental Status Exam; MNA-SF—Mini Nutritional Assessment—Short Form; MOS—Medical Outcomes Study; NHL—non-Hodgkin Lymphoma; OS—overall survival; PHQ-9—Patient Health Questionnaire-9; R-MCI—revised Myeloma Comorbidity Index; SIOG—International Society of Geriatric Oncology; VES-13—Vulnerable Elders Survey-13

Note. Independent indicates that the tool is used as a known single tool for frailty; combinatorial indicates that the tool is used as a combination of tools, with frailty assessed as a part of that.

as geriatric assessments; however, there are variations in components that constitute these geriatric assessments. Some include functional status, comorbidities, and age, whereas others include additional assessments of cognition and psychosocial dysfunction (Engelhardt et al., 2016; Ferrat et al., 2017; Marchesi et al., 2013; Palumbo et al., 2015; Velghe et al., 2016; Zhong et al., 2017). The time-intensive nature of the CGA is probably one of the reasons for studies using selective, fewer, and shorter assessments for geriatric assessment, instead of the more time-consuming and extensive measures in the CGA. Some of the tools used as a part of geriatric assessments include the VES-13, G8, and Karnofsky Performance Status (KPS) scale.

Tools such as the VES-13 and G8 have been analyzed to determine their respective predictive ability for abnormal geriatric assessment or frailty. The VES-13 is a 13-item survey that includes age, as well as self-rated health and functional status, and is scored from 0 to 10, with higher scores indicating worse or more extensive

frailty (Saliba et al., 2001). The G8 consists of eight domains: food intake, weight loss, body mass index, mobility, neuropsychological problems, prescription drugs, health status, and age. In recipients of allogeneic HSCT, the G8 correlates with the CHS frailty index and showed a higher sensitivity (69.7%), whereas the VES-13 had higher specificity (100%) (Holmes et al., 2014).

A study by Muffly et al. (2013) showed that 83% of patients who had performance status scores indicating good health fell into either the frail or pre-frail category based on the CHS frailty index (incorporated as part of the CGA). Similarly, KPS allows patients to be classified based on their functional impairment; it also demonstrates that patients can have a KPS score of greater than 80 (indicating good functional status) but also have an abnormal geriatric assessment (Crooks et al., 1991; Holmes et al., 2014). These studies clearly indicate that individual screening tools (e.g., World Health Organization Performance Status, KPS) may not be

TABLE 5. Assessment Tools Used in Selected Studies

Tool	Description	Articles Using Tool
ADLs	Questionnaire that assesses ability to perform basic daily activities	Corsetti et al., 2011; Engelhardt et al., 2016; Ferrat et al., 2017; Holmes et al., 2014; Marchesi et al., 2013; Milani et al., 2016; Molica et al., 2019; Muffly et al., 2013, 2014; Okuyama et al., 2015; Palumbo et al., 2015; Velghe et al., 2016; Zhong et al., 2017
Balducci geriatric assessment	Based on age, ADLs, IADLs, CIRS-G, presence of geriatric syndrome	Ferrat et al., 2017
CCI	Tool used to assess comorbidity by assigning weights to chronic conditions	Buckstein et al., 2016; Budziszewska et al., 2015; Engelhardt et al., 2016, 2017; Hshieh et al., 2018; Muffly et al., 2013; Offidani et al., 2012; Palumbo et al., 2015; Sakatoku et al., 2019; Zhong et al., 2017
CFS	7- or 9-point visual scale that serves as a clinical measure of fitness and frailty	Buckstein et al., 2016; Hshieh et al., 2018; Sakatoku et al., 2019
CGA	Interprofessional tool that considers psychosocial, cognitive, financial, environmental, and functional limitations	Corsetti et al., 2011; Marchesi et al., 2013; Muffly et al., 2013, 2014; Velghe et al., 2016
CHS frailty index	Phenotypic assessment of frailty that includes decreased grip strength, decreased gait speed, decreased physical activity, weight loss, and fatigue	Arora et al., 2016; Eissa et al., 2017; Engelhardt et al., 2016, 2017; Holmes et al., 2014; Hshieh et al., 2018; Muffly et al., 2013, 2014; Wilson et al., 2016
CIRS	Scale that summarizes overall severity of illness based on clinical information	Corsetti et al., 2011; Ferrat et al., 2017; Marchesi et al., 2013; Molica et al., 2019; Muffly et al., 2013
ECOG-PS	Describes level of functioning related to ability to care for oneself, daily activity, and physical activity	Buckstein et al., 2016; Budziszewska et al., 2015; Ferrat et al., 2017; Muffly et al., 2013, 2014
G8	Screening tool that includes food intake, weight loss, mobility, neuropsychological problems, body mass index, prescription drugs, health status, and age; 8 domains	Holmes et al., 2014; Velghe et al., 2016
GFI	15-item questionnaire that assesses physical, cognitive, social, and psychological domains	Park et al., 2015
HCT-CI	Comorbidity index that comprises 17 categories of organ dysfunction	Engelhardt et al., 2016, 2017; Holmes et al., 2014; Muffly et al., 2013, 2014; Rosko et al., 2019
HGS	Basic measure to determine musculoskeletal function	Eissa et al., 2017; Muffly et al., 2013, 2014; Velghe et al., 2016; Wilson et al., 2016
IADLs	Questionnaire that assesses ability to independently perform basic activities and functional skills	Buckstein et al., 2016; Corsetti et al., 2011; Engelhardt et al., 2016; Ferrat et al., 2017; Holmes et al., 2014; Milani et al., 2016; Molica et al., 2019; Muffly et al., 2013, 2014; Okuyama et al., 2015; Palumbo et al., 2015; Velghe et al., 2016; Zhong et al., 2017
KFI	Index focusing on type of comorbidity and pathophysiologic severity of comorbid conditions; classified as vascular or nonvascular diseases	Engelhardt et al., 2016, 2017
KPS	Index that allows for classification based on functional impairment	Rosko et al., 2019

Continued on the next page

TABLE 5. Assessment Tools Used in Selected Studies (Continued)

Tool	Description	Articles Using Tool
LC typology geriatric assessment	Inadequate social environment, malnutrition, depression, cognitive impairment, CIRS-G, ADLs, age, and tumor size	Ferrat et al., 2017
SIOG 1	Based on ADLs, IADLs, CIRS-G, malnutrition based on weight loss, and ECOG-PS	Ferrat et al., 2017
SIOG 2	2-step process—if abnormal G8, proceed to step 2; includes ADLs, IADLs, CIRS-G, malnutrition based on weight loss, and neuropsychological problems	Ferrat et al., 2017
VES-13	Questionnaire about age, self-rated health, and functional status	Holmes et al., 2014; Okuyama et al., 2015
Vulnerability score	Assessment that includes age, performance status, renal insufficiency, bone fracture, cytopenia, and comorbidity	Offidani et al., 2012

ADLs—activities of daily living; CCI—Charlson Comorbidity Index; CFS—Clinical Frailty Scale; CGA—Comprehensive Geriatric Assessment; CHS—Cardiovascular Health Study; CIRS—Cumulative Illness Rating Scale; CIRS-G—Cumulative Illness Rating Scale—Geriatric; ECOG-PS—Eastern Cooperative Oncology Group Performance Status; GFI—Groningen Frailty Index; HCT-CI—Hematopoietic Cell Transplantation-Specific Comorbidity Index; HGS—hand grip strength; IADLs—instrumental activities of daily living; KFI—Kaplan-Feinstein Index; KPS—Karnofsky Performance Status; SIOG—International Society of Geriatric Oncology; VES-13—Vulnerable Elders Survey-13

able to comprehensively assess frailty and overall health.

These results also show that although shorter geriatric assessment tools are available, they do have certain drawbacks. Using a consistent combination of tools may be important to accurately determine comorbidities, vulnerability, and frailty. It is important to acknowledge these discrepancies and come to a uniform consensus on shorter tools to be used in this specific population, if needed. At this time, there is a lack of evidence to strongly recommend a relatively shorter assessment tool that is sensitive and specific.

Frailty and Comorbidities

Although frailty and comorbidities coexist, they have been identified as distinct entities. Measures of frailty often include the presence of comorbidities, potentially hindering an absolute measure of frailty. Comorbidity has been defined as any distinct additional health condition that has existed or may occur during the clinical course of a patient with a primary disease (Feinstein, 1970). Some studies in patients undergoing HSCT have adopted a combinatorial approach involving performance status, frailty, and comorbidity assessments for stratifying at-risk patients. The Charlson Comorbidity Index (CCI) is calculated by assigning weights to chronic conditions; this tool has been widely used in various chronic

conditions (Charlson et al., 1987; Extermann, 2000). Offidani et al. (2012) conducted a study involving a vulnerability score that combined both performance status and the CCI. This score proved to better predict the survival of patients with multiple myeloma versus a comparison of patients using performance status or the CCI alone. Importantly, Offidani et al.'s (2012) results suggest that this vulnerability score can identify two key groups of patients: (a) those who can tolerate effective therapy and (b) those who may experience nonmyeloma-related mortality, indicating great promise for risk stratification.

The Myelodysplastic Syndrome-Specific Comorbidity Index (MDS-CI) is another dynamic index that is used to predict the outcome of patients with myelodysplastic syndrome (Buckstein et al., 2016). In a study by Buckstein et al. (2016), the predictive value of a prognosis scoring system improved when frailty and comorbidity measures were added. Adding the CFS and the CCI/MDS-CI comorbidity tools increased the predictive value of the scoring system by 30% and 5%, respectively, leading to a more precise prediction of overall survival (Buckstein et al., 2016; Sakatoku et al., 2019).

With the intention of establishing comorbidity scores in patients undergoing HSCT, Sorror et al. (2005) introduced a new simple index, the Hematopoietic Cell Transplantation-Specific Comorbidity Index (HCT-CI), that provides valid

and reliable scoring for patients pretransplantation. HCT-CI scores consider 17 categories of organ dysfunction that have the potential to predict nonrelapse mortality and survival. This tool is used in HSCT pre-assessments (Sorrer et al., 2005). The HCT-CI has been incorporated within geriatric assessments and in combination with functional status assessments, and has served as an improved stratification tool (Engelhardt et al., 2017; Holmes et al., 2014; Muffy et al., 2013, 2014; Rosko et al., 2019).

In summary, comorbidity tools (e.g., CCI, MDS-CI, HCT-CI) emphasize the importance of assessing patients for comorbidities but also affirm that the concomitant use of independent frailty assessments (e.g., performance status, CFS, vulnerability score) add to eligibility screening, stratification of at-risk patients, and the predictive value of outcomes. However, in carrying out future research, it is essential that there be agreement on how both comorbidities and frailty are defined and assessed.

Although this article primarily focuses on frailty assessment in adults undergoing HSCT, it is important to acknowledge the incidence of frailty in childhood survivors of HSCT. A study among adult survivors of childhood hematologic cancers demonstrated a four-fold increase in the incidence of frailty using the CHS frailty index in recipients of HSCT compared to survivors treated with conventional therapy (Eissa et al., 2017). Overall, there is a need to assess frailty, identify the appropriate tool, and demonstrate the value of combining tools for frailty and comorbidity in this patient population.

Frailty in Patients With Hematologic Malignancies

During the study selection process for this scoping review, the current authors identified cohort studies that addressed frailty in hematologic malignancies and listed “HSCT” in the article abstract or body and, therefore, met the inclusion criteria for this review. Of the 13 studies grouped under hematologic malignancies, 3 involved patients with a diagnosis of multiple myeloma and 2 were in patients being treated for lymphomas; the remaining studies looked at patients with leukemias and other hematologic cancers. The current authors reviewed tools that were used to assess frailty among patients with hematologic malignancies. Some studies proposed the use of prescreening tools that measure features of frailty, such as hand grip strength and the Mini Nutritional Assessment–Short Form (Park et al., 2015; Velghe et al., 2016). In a study by Velghe et al. (2016), decreased hand grip strength showed significant association with

KNOWLEDGE TRANSLATION

- Frailty assessment should be incorporated for risk stratification of individuals with hematologic malignancies and recipients of hematopoietic stem cell transplantation.
 - Evidence is lacking to identify the “best” frailty assessment tool.
 - Combinatorial assessment tools may have more predictive power than independent assessment tools.
-

abnormal G8 scores in women but not in men. These prescreening tools show associations with abnormal geriatric assessments and have a predictive ability of overall survival in these patients. In some instances, tools such as the CHS frailty index and CFS were implemented in patients with leukemia and myeloma, showing associations of frailty with outcomes such as decreased bone mass density and impaired cognitive function (Hshieh et al., 2018; Wilson et al., 2016). There were several studies reviewed for the current article that indicated the full use of the CGA; however, further exploration revealed that only select assessments had been used as part of the overall CGA (e.g., comorbidity, functional status, incontinence, falls), whereas other CGA domains were not measured. Therefore, it was difficult to interpret the utility of the CGA equally across the studies reviewed.

Use of Combinatorial Assessments

Building on individual tools, researchers have explored and initiated a combinatorial approach for assessing frailty in hematologic malignancies, similar to that performed in recipients of HSCT, as previously discussed. Review of the hematologic cancer literature showed predominant use of functional assessment, such as activities of daily living (ADLs) and instrumental ADLs (IADLs); age; and comorbidity as measures of frailty in blood cancer diagnosis (Engelhardt et al., 2016; Mellqvist, 2015). IMWG score combined age, ADLs, IADLs, and the CCI and was able to stratify based on fitness and predict the risk of mortality in patients newly diagnosed with multiple myeloma (Palumbo et al., 2015). Similar studies evaluated use of the VES-13, Eastern Cooperative Oncology Group Performance Status, the CCI, and age for stratifying individuals with high-risk blood cancer (Budziszewska et al., 2015; Milani et al., 2016; Okuyama et al., 2015). Combined use of the VES-13 and assessment of anhedonia among older patients with newly diagnosed cancer showed improved sensitivity and negative predictive value of frailty (Okuyama et al., 2015). Although it is understandable that each cancer type has

its own assessment, the data provide compelling evidence to believe that combining assessment tools will help to improve stratification of high-risk patients, identification of patients for managing treatment toxicities, and the predictive power of outcomes (e.g., overall survival, disease-free survival, mortality) in these patients (Corsetti et al., 2011; Marchesi et al., 2013; Molica et al., 2019; Spina et al., 2012).

Discussion

Healthcare providers have used chronological age, various clinical parameters, and different assessments to determine eligibility for specific treatment regimens (e.g., chemotherapy, HSCT) for patients, as well as prognosis and outcomes. With the advent of novel therapeutic agents and reduced-intensity chemotherapy regimens, there has been an increase in the number of patients who are being treated; consequently, providers are increasingly being presented with the challenge of accurately determining eligibility for various treatments and procedures. Clinicians and scientists have stated repeatedly that there is a need to improve and refine existing preassessments (Tay et al., 2019). Frailty has been associated with increased toxicity, poor therapeutic response, and decreased survival for patients with hematologic malignancies (Ness et al., 2013; Sorrow et al., 2005). Research shows that pre-HSCT prevalence of frailty is higher in patients requiring HSCT compared to the general geriatric population (Artz et al., 2011; Muffly et al., 2014). As the average lifespan increases, healthcare providers will encounter more patients who are aging and frail, emphasizing the need to have an established plan for stratifying patients who will be eligible for HSCT (Hamaker et al., 2012; Mohile & Magnuson, 2013). Such assessments will help in identifying those who will be able to tolerate transplantation, improve informed decision-making regarding therapy modality, and monitor response and overall quality of life by providing appropriate supportive care (Holmes et al., 2014).

The purpose of this scoping review was to systematically gather data on how frailty has been assessed in recipients of HSCT and patients with hematologic malignancies. Tools such as the CHS frailty index, CFS, VES-13, hand grip strength, G8, and CGA have been used in various settings in patients undergoing HSCT. Inconsistencies in how frailty is defined and assessed present as a barrier in applying data to larger cohorts of patients and providing guidelines. The current synthesis of existing literature demonstrates that frailty has been assessed independently using established frailty tools such as the CHS frailty

index, surrogate measures such as hand grip strength, combinatorial assessments (which typically include a combination of performance status, functional status, and comorbidities), and the time- and personnel-intensive CGA.

In addition, the current authors encountered variable measures that present challenges for quantitatively evaluating and recommending tools for implementation. For example, in studies using the CHS frailty index, the physical activity component was either a self-reported question or a compendium of physical activities using metabolic rates. With geriatric assessments, the individual components that comprise the physical activity score and the evaluation methods or tools used were varied. Such variability in assessing a component of a measure makes it challenging to compare data across various clinical settings or studies. The task at hand is to determine the tool, or combination of tools, that should be used to assess frailty, with specific instructions on how to administer the assessment in a uniform manner.

Research has repeatedly shown the benefits of combining assessment tools for frailty and comorbidity to help improve healthcare providers' ability to stratify high-risk patients, which also improves patient outcomes, such as overall survival. An array of comorbidity assessment tools are used in patients with hematologic malignancies or in recipients of HSCT (CCI, R-MCI, HCT-CI). Although these three tools are all broadly considered comorbidity indices, the CCI specifically measures chronic conditions, the HCT-CI measures organ dysfunction, and the R-MCI measures frailty (via the CHS frailty index) in addition to organ dysfunction, contributing to the overall scores. Given the compelling evidence in favor of combining tools to arrive at a more holistic and accurate measure of frailty, the current authors suggest the testing and implementation of a frailty and comorbidity assessment tool to evaluate patients, but also recommend that clinicians and researchers adhere to the same tools and combination of tools for their specific patient populations. For example, the R-MCI could be used for all patients with myeloma, whereas a combination of the HCT-CI and CHS frailty index could be used for all patients undergoing HSCT, regardless of the underlying diagnosis. Such recommendations would serve as an important first step to better stratify patients that may, ultimately, lead to the identification of patient subgroups that could benefit from planned interventions. The Composite

Health Assessment Risk Model (CHARM) study conducted by the Blood and Marrow Transplant Clinical Trials Network (BMT CTN 1704) plans to investigate a composite health assessment model that includes comorbidity and geriatric assessment in older adults receiving allogeneic HSCT (Artz & Sorror, 2021). Results from this study have the potential to provide information that could help clinicians and researchers shape their preassessment strategies.

Frailty should be incorporated as part of preassessment and risk stratification for potential recipients of HSCT and patients with hematologic malignancies. The CHS frailty index is most commonly used across studies, but the combination of assessment tools (e.g., concurrent use of a frailty tool and comorbidity index) has been shown to have more predictive power. The specific tools that will contribute to the best cumulative frailty score remain to be determined; however, once identified, all researchers and clinicians should adhere to predetermined assessment tools to maintain consistency, which will enable early implementation of supportive therapy in this patient population. These strategies will better inform personalized treatment planning and supportive interventions (early implementation of physical therapy, occupational therapy, lifestyle counseling, and dietary guidance), as necessary (Chemaitilly & Robison, 2012; Hegde & Murthy, 2018).

Implications for Nursing

Healthcare providers, like advanced nurse practitioners (ANPs) and nurses, play a significant role in providing patient-centered care, which involves prevention and risk assessment in patients with cancer (Morgan & Tarbi, 2016). This is also applicable in patients slated to receive HSCT, recipients of HSCT, and patients diagnosed with hematologic cancers. ANPs and nurses are involved in the initial consultation with these patients; this involves determining baseline health history information, diagnosis, comorbidities, barriers and challenges (physical and emotional), and caregiver support. They are well positioned to conduct frailty assessments; in conjunction with the interprofessional transplantation team, they can help to facilitate appropriate treatment planning and provide personalized care. The involvement of APNs and nurses throughout the care continuum, established trust, effective communication, and collaboration with the transplantation team give nurses the unique ability to lead the way in assessing and addressing frailty as an essential component of treatment planning and individualized interventions in recipients of HSCT and patients

diagnosed with hematologic malignancies. Based on information gathered in this scoping review, it is recommended that nurse clinicians and researchers incorporate frailty assessments as a preassessment and a risk-stratification tool, as well as conduct clinical and translational research to determine appropriate frailty assessment tools. This should be followed by consistent administration of the identified tools to assess frailty in this population.

Conclusion

Frailty is an important parameter for understanding an individual's "physiological age" and developing personalized treatment plans for patients with hematologic malignancies and recipients of HSCT. Frailty assessments are being used by healthcare providers; however, there is a need for consistency on how frailty is defined and assessed. A combinatorial approach that includes evaluation of frailty and comorbidity has shown improved predictive power of patient outcomes. The first step is to identify the tools of choice and administer them in a consistent manner. This can subsequently lead to the initiation and implementation of appropriate interventions in a timely manner that will help to improve and sustain a better quality of life in these patient populations.

Lathika Mohanraj, PhD, RN, BMTCN[®], and **Lana Sargent, PhD, RN, FNP-C, GNP-BC**, are both assistant professors in the School of Nursing, **Roy Brown, MLIS, AHIP**, is an associate professor, interim co-department head of the Research and Education Department, research and education librarian in the Tompkins-McCaw Library for the Health Sciences, and liaison to the School of Nursing, and **Theresa Swift-Scanlan, PhD, RN, FAAN**, is an associate professor and the director of the Biobehavioral Research Laboratory in the School of Nursing, all at the Virginia Commonwealth University in Richmond. Mohanraj can be reached at mohanraj@vcu.edu, with copy to ONFEditor@ons.org. (Submitted August 2020. Accepted October 10, 2020.)

This research was supported by the Virginia Commonwealth University Presidential Research Quest Fund, awarded to Mohanraj.

All authors contributed to the conceptualization and design, completed the data collection, provided the analysis, and contributed to the manuscript preparation.

REFERENCES

- Abel, G.A., & Klepin, H.D. (2018). Frailty and the management of hematologic malignancies. *Blood*, 131(5), 515–524. <https://doi.org/10.1182/blood-2017-09-746420>

- Arora, M., Sun, C.-L., Ness, K.K., Teh, J.B., Wu, J., Francisco, L., . . . Bhatia, S. (2016). Physiologic frailty in nonelderly hematopoietic cell transplantation patients: Results from the bone marrow transplant survivor study. *JAMA Oncology*, 2(10), 1277–1286. <https://doi.org/10.1001/jamaoncol.2016.0855>
- Artz, A., & Sorror, M.L. (2021, March 11). Composite Health Assessment Risk Model (CHARM) for older adults (BMT CTN 1704). *ClinicalTrials.gov*. <https://clinicaltrials.gov/ct2/show/NCT03992352>
- Artz, A., Swanson, K., Kocherginsky, M., Godley, L., Kline, J., Nguyen, V., . . . van Besien, K. (2011). Features of frailty are surprisingly common in adults 50 years and older undergoing allogeneic hematopoietic cell transplantation (HCT) in the modern era [Abstract 409]. *Biology of Blood and Marrow Transplantation*, 17(2, Suppl.), S302. <https://doi.org/10.1016/j.bbmt.2010.12.445>
- Buckstein, R., Wells, R.A., Zhu, N., Leitch, H.A., Nevill, T.J., Yee, K.W.L., . . . Alibhai, S.M.H. (2016). Patient-related factors independently impact overall survival in patients with myelodysplastic syndromes: An MDS-CAN prospective study. *British Journal of Haematology*, 174(1), 88–101. <https://doi.org/10.1111/bjh.14033>
- Budziszewska, B.K., Pluta, A., Sulek, K., Wierzbowska, A., Robak, T., Giebel, S., . . . Lech-Maranda, E. (2015). Treatment of elderly patients with acute myeloid leukemia adjusted for performance status and presence of comorbidities: A Polish Adult Leukemia Group study. *Leukemia and Lymphoma*, 56(8), 2331–2338. <https://doi.org/10.3109/10428194.2014.985672>
- Charlson, M.E., Pompei, P., Ales, K.L., & MacKenzie, C.R. (1987). A new method of classifying prognostic comorbidity in longitudinal studies: Development and validation. *Journal of Chronic Diseases*, 40(5), 373–383. [https://doi.org/10.1016/0021-9681\(87\)90171-8](https://doi.org/10.1016/0021-9681(87)90171-8)
- Chemaitilly, W., & Robison, L.L. (2012). Safety of growth hormone treatment in patients previously treated for cancer. *Endocrinology and Metabolism Clinics of North America*, 41(4), 785–792. <https://doi.org/10.1016/j.ecl.2012.07.002>
- Corsetti, M.T., Salvi, F., Perticone, S., Baraldi, A., De Paoli, L., Gatto, S., . . . Levis, A. (2011). Hematologic improvement and response in elderly AML/RAEB patients treated with valproic acid and low-dose Ara-C. *Leukemia Research*, 35(8), 991–997. <https://doi.org/10.1016/j.leukres.2011.02.021>
- Crooks, V., Waller, S., Smith, T., & Hahn, T.J. (1991). The use of the Karnofsky Performance Scale in determining outcomes and risk in geriatric outpatients. *Journal of Gerontology*, 46(4), M139–M144. <https://doi.org/10.1093/geronj/46.4.M139>
- Domm, A.-S., Hieke, S., Ihorst, G., Pantic, M., Koch, B., Kleber, M., . . . Engelhardt, M. (2014). Importance and determinants of comorbidities, functional limitations and multiple myeloma (MM)-specific risk factors: Further development of an improved and weighted MM-risk score (Freiburg Comorbidity Index [FCI]). *Blood*, 124(21), 733. <https://doi.org/10.1182/blood.v124.21.733-733>
- Eissa, H.M., Lu, L., Baassiri, M., Bhakta, N., Ehrhardt, M.J., Triplett, B.M., . . . Ness, K.K. (2017). Chronic disease burden and frailty in survivors of childhood HSCT: A report from the St. Jude Lifetime Cohort Study. *Blood Advances*, 1(24), 2243–2246. <https://doi.org/10.1182/bloodadvances.2017010280>
- El Assar, M., Laosa, O., & Rodríguez Mañas, L. (2019). Diabetes and frailty. *Current Opinion in Clinical Nutrition and Metabolic Care*, 22(1), 52–57. <https://doi.org/10.1097/MCO.0000000000000535>
- Engelhardt, M., Dold, S.M., Ihorst, G., Zober, A., Möller, M., Reinhardt, H., . . . Wäsch, R. (2016). Geriatric assessment in multiple myeloma patients: Validation of the International Myeloma Working Group (IMWG) score and comparison with other common comorbidity scores. *Haematologica*, 101(9), 1110–1119. <https://doi.org/10.3324/haematol.2016.148189>
- Engelhardt, M., Domm, A.-S., Dold, S.M., Ihorst, G., Reinhardt, H., Zober, A., . . . Wäsch, R. (2017). A concise revised Myeloma Comorbidity Index as a valid prognostic instrument in a large cohort of 801 multiple myeloma patients. *Haematologica*, 102(5), 910–921. <https://doi.org/10.3324/haematol.2016.162693>
- Extermann, M. (2000). Measuring comorbidity in older cancer patients. *European Journal of Cancer*, 36(4), 453–471.
- Extermann, M., & Hurria, A. (2007). Comprehensive geriatric assessment for older patients with cancer. *Journal of Clinical Oncology*, 25(14), 1824–1831. <https://doi.org/10.1200/JCO.2007.10.6559>
- Feinstein, A.R. (1970). The pre-therapeutic classification of co-morbidity in chronic disease. *Journal of Chronic Diseases*, 23(7), 455–468. [https://doi.org/10.1016/0021-9681\(70\)90054-8](https://doi.org/10.1016/0021-9681(70)90054-8)
- Ferrat, E., Paillaud, E., Caillet, P., Laurent, M., Tournigand, C., Lagrange, J.-L., . . . Bastuji-Garin, S. (2017). Performance of four frailty classifications in older patients with cancer: Prospective elderly cancer patients cohort study. *Journal of Clinical Oncology*, 35(7), 766–777. <https://doi.org/10.1200/JCO.2016.69.3143>
- Fried, L.P., Tangen, C.M., Walston, J., Newman, A.B., Hirsch, C., Gottdiener, J., . . . McBurnie, M.A. (2001). Frailty in older adults: Evidence for a phenotype. *Journals of Gerontology Series A*, 56(3), M146–M157. <https://doi.org/10.1093/gerona/56.3.m146>
- Hamaker, M.E., Jonker, J.M., de Rooij, S.E., Vos, A.G., Smorenburg, C.H., & van Munster, B.C. (2012). Frailty screening methods for predicting outcome of a comprehensive geriatric assessment in elderly patients with cancer: A systematic review. *Lancet Oncology*, 13(10), E437–E444. [https://doi.org/10.1016/S1470-2045\(12\)70259-0](https://doi.org/10.1016/S1470-2045(12)70259-0)
- Handforth, C., Clegg, A., Young, C., Simpkins, S., Seymour, M.T., Selby, P.J., & Young, J. (2015). The prevalence and outcomes of frailty in older cancer patients: A systematic review. *Annals of Oncology*, 26(6), 1091–1101. <https://doi.org/10.1093/annonc/mdu540>
- Hegde, A., & Murthy, H.S. (2018). Frailty: The missing piece of the pre-hematopoietic cell transplantation assessment? *Bone*

- Marrow Transplantation*, 53(1), 3–10. <https://doi.org/10.1038/bmt.2017.192>
- Holmes, H.M., Des Bordes, J.K.A., Kebriaei, P., Yennu, S., Champlin, R.E., Giralt, S., & Mohile, S.G. (2014). Optimal screening for geriatric assessment in older allogeneic hematopoietic cell transplantation candidates. *Journal of Geriatric Oncology*, 5(4), 422–430. <https://doi.org/10.1016/j.jgo.2014.04.004>
- Hsieh, T.T., Jung, W.F., Grande, L.J., Chen, J., Stone, R.M., Soiffer, R.J., . . . Abel, G.A. (2018). Prevalence of cognitive impairment and association with survival among older patients with hematologic cancers. *JAMA Oncology*, 4(5), 686–693. <https://doi.org/10.1001/jamaoncol.2017.5674>
- Marchesi, F., Cenfra, N., Altomare, L., Dessanti, M.L., Mecarocci, S., Cerchiara, E., . . . Cimino, G. (2013). A retrospective study on 73 elderly patients (≥ 75 years) with aggressive B-cell non Hodgkin lymphoma: Clinical significance of treatment intensity and comprehensive geriatric assessment. *Journal of Geriatric Oncology*, 4(3), 242–248. <https://doi.org/10.1016/j.jgo.2013.04.003>
- Mellqvist, U.-H. (2015). New prognostic tools for myeloma. *Blood*, 125(13), 2014–2015. <https://doi.org/10.1182/blood-2015-02-626408>
- Milani, P., Rajkumar, S.V., Merlini, G., Kumar, S., Gertz, M.A., Palladini, G., . . . Dispenzieri, A. (2016). N-terminal fragment of the type-B natriuretic peptide (NT-proBNP) contributes to a simple new frailty score in patients with newly diagnosed multiple myeloma. *American Journal of Hematology*, 91(11), 1129–1134. <https://doi.org/10.1002/ajh.24532>
- Mohile, S.G., & Magnuson, A. (2013). Comprehensive geriatric assessment in oncology. *Interdisciplinary Topics in Gerontology*, 38, 85–103. <https://doi.org/10.1159/000343608>
- Molica, S., Giannarelli, D., Levato, L., Mirabelli, R., Levato, D., Lentini, M., & Piro, E. (2019). A simple score based on geriatric assessment predicts survival in elderly newly diagnosed chronic lymphocytic leukemia patients. *Leukemia and Lymphoma*, 60(3), 845–847. <https://doi.org/10.1080/10428194.2018.1508674>
- Morgan, B., & Tarbi, E. (2016). The role of the advanced practice nurse in geriatric oncology care. *Seminars in Oncology Nursing*, 32(1), 33–43. <https://doi.org/10.1016/j.soncn.2015.11.005>
- Morley, J.E., Vellas, B., Abellan van Kan, G., Anker, S.D., Bauer, J.M., Bernabei, R., . . . Walston, J. (2013). Frailty consensus: A call to action. *Journal of the American Medical Directors Association*, 14(6), 392–397. <https://doi.org/10.1016/j.jamda.2013.03.022>
- Morrison, V.A., Hamlin, P., Soubeyran, P., Stauder, R., Wadhwa, P., Aapro, M., & Lichtman, S. (2015). Diffuse large B-cell lymphoma in the elderly: Impact of prognosis, comorbidities, geriatric assessment, and supportive care on clinical practice. An International Society of Geriatric Oncology (SIOG) expert position paper. *Journal of Geriatric Oncology*, 6(2), 141–152. <https://doi.org/10.1016/j.jgo.2014.11.004>
- Muffy, L.S., Boulukos, M., Swanson, K., Kocherginsky, M., del Cerro, P., Schroeder, L., . . . Artz, A.S. (2013). Pilot study of comprehensive geriatric assessment (CGA) in allogeneic transplant: CGA captures a high prevalence of vulnerabilities in older transplant recipients. *Biology of Blood and Marrow Transplantation*, 19(3), 429–434. <https://doi.org/10.1016/j.bbmt.2012.11.006>
- Muffy, L.S., Kocherginsky, M., Stock, W., Chu, Q., Bishop, M.R., Godley, L.A., . . . Artz, A.S. (2014). Geriatric assessment to predict survival in older allogeneic hematopoietic cell transplantation recipients. *Haematologica*, 99(8), 1373–1379. <https://doi.org/10.3324/haematol.2014.103655>
- Ness, K.K., Krull, K.R., Jones, K.E., Mulrooney, D.A., Armstrong, G.T., Green, D.M., . . . Hudson, M.M. (2013). Physiologic frailty as a sign of accelerated aging among adult survivors of childhood cancer: A report from the St Jude Lifetime Cohort Study. *Journal of Clinical Oncology*, 31(36), 4496–4503. <https://doi.org/10.1200/JCO.2013.52.2268>
- Offidani, M., Corvatta, L., Polloni, C., Centurioni, R., Visani, G., Brunori, M., . . . Leoni, P. (2012). Assessment of vulnerability measures and their effect on survival in a real-life population of multiple myeloma patients registered at Marche Region Multiple Myeloma Registry. *Clinical Lymphoma, Myeloma and Leukemia*, 12(6), 423–432. <https://doi.org/10.1016/j.clml.2012.06.008>
- Okuyama, T., Sugano, K., Iida, S., Ishida, T., Kusumoto, S., & Akechi, T. (2015). Screening performance for frailty among older patients with cancer: A cross-sectional observational study of two approaches. *Journal of the National Comprehensive Cancer Network*, 13(12), 1525–1531. <https://doi.org/10.6004/jnccn.2015.0180>
- Palumbo, A., Bringhen, S., Mateos, M.-V., Larocca, A., Facon, T., Kumar, S.K., . . . Rajkumar, S.V. (2015). Geriatric assessment predicts survival and toxicities in elderly myeloma patients: An International Myeloma Working Group report. *Blood*, 125(13), 2068–2074. <https://doi.org/10.1182/blood-2014-12-615187>
- Park, S., Hong, J., Hwang, I., Ahn, J.-Y., Cho, E.Y., Park, J., . . . Lee, J.H. (2015). Comprehensive geriatric assessment in elderly patients with newly diagnosed aggressive non-Hodgkin lymphoma treated with multi-agent chemotherapy. *Journal of Geriatric Oncology*, 6(6), 470–478. <https://doi.org/10.1016/j.jgo.2015.10.183>
- Pel-Littel, R.E., Schuurmans, M.J., Emmelot-Vonk, M.H., & Verhaar, H.J.J. (2009). Frailty: Defining and measuring of a concept. *Journal of Nutrition, Health and Aging*, 13(4), 390–394. <https://doi.org/10.1007/s12603-009-0051-8>
- Perna, S., Francis, M.D.A., Bologna, C., Moncaglieri, F., Riva, A., Morazzoni, P., . . . Rondanelli, M. (2017). Performance of Edmonton Frail Scale on frailty assessment: Its association with multi-dimensional geriatric conditions assessed with specific screening tools. *BMC Geriatrics*, 17, 2. <https://doi.org/10.1186/s12877-016-0382-3>

- Phelan, R., Arora, M., & Chen, M. (2020). Current use and outcome of hematopoietic stem cell transplantation. Center for International Blood and Marrow Transplant Research. <https://www.cibmtr.org/ReferenceCenter/SlidesReports/SummarySlides/pages/index.aspx>
- Pidala, J., Craig, B.M., Lee, S.J., Majhail, N., Quinn, G., & Anasetti, C. (2013). Practice variation in physician referral for allogeneic hematopoietic cell transplantation. *Bone Marrow Transplantation*, 48(1), 63–67. <https://doi.org/10.1038/bmt.2012.95>
- Rosko, A., Hofmeister, C., Benson, D., Efebera, Y., Huang, Y., Gillahan, J., . . . Burd, C.E. (2015). Autologous hematopoietic stem cell transplant induces the molecular aging of T-cells in multiple myeloma. *Bone Marrow Transplantation*, 50(10), 1379–1381. <https://doi.org/10.1038/bmt.2015.143>
- Rosko, A.E., Huang, Y., Benson, D.M., Efebera, Y.A., Hofmeister, C., Jaglowski, S., . . . Burd, C.E. (2019). Use of a comprehensive frailty assessment to predict morbidity in patients with multiple myeloma undergoing transplant. *Journal of Geriatric Oncology*, 10(3), 479–485. <https://doi.org/10.1016/j.jgo.2018.05.015>
- Sakataku, K., Takeoka, Y., Miura, A., Araki, T., Fujitani, Y., Yamamura, R., . . . Hino, M. (2019). Combination of frailty status and comorbidity score improves the stratification of survival in patients with myelodysplastic syndrome owing to good predictive capability for infection-related mortality. *Clinical Lymphoma, Myeloma and Leukemia*, 19(12), 799–805. <https://doi.org/10.1016/j.clml.2019.09.610>
- Saliba, D., Elliott, M., Rubenstein, L.Z., Solomon, D.H., Young, R.T., Kamberg, C.J., . . . Wenger, N.S. (2001). The Vulnerable Elders Survey: A tool for identifying vulnerable older people in the community. *Journal of the American Geriatrics Society*, 49(12), 1691–1699. <https://doi.org/10.1046/j.1532-5415.2001.49281.x>
- Searle, S.D., & Rockwood, K. (2015). Frailty and the risk of cognitive impairment. *Alzheimer's Research and Therapy*, 7(1), 54. <https://doi.org/10.1186/s13195-015-0140-3>
- Song, X., Mitnitski, A., & Rockwood, K. (2014). Age-related deficit accumulation and the risk of late-life dementia. *Alzheimer's Research and Therapy*, 6(5), 54. <https://doi.org/10.1186/s13195-014-0054-5>
- Sorrow, M.L., Maris, M.B., Storb, R., Baron, F., Sandmaier, B.M., Maloney, D.G., & Storer, B. (2005). Hematopoietic Cell Transplantation (HCT)-Specific Comorbidity Index: A new tool for risk assessment before allogeneic HCT. *Blood*, 106(8), 2912–2919. <https://doi.org/10.1182/blood-2005-05-2004>
- Spina, M., Balzarotti, M., Uziel, L., Ferreri, A.J.M., Fratino, L., Magagnoli, M., . . . Tirelli, U. (2012). Modulated chemotherapy according to modified comprehensive geriatric assessment in 100 consecutive elderly patients with diffuse large B-cell lymphoma. *Oncologist*, 17(6), 838–846. <https://doi.org/10.1634/theoncologist.2011-0417>
- Tay, J., Daly, A., Jamani, K., Labelle, L., Savoie, L., Stewart, D., . . . Beattie, S. (2019). Patient eligibility for hematopoietic stem cell transplantation: A review of patient-associated variables. *Bone Marrow Transplantation*, 54(3), 368–382.
- Tricco, A.C., Lillie, E., Zarin, W., O'Brien, K.K., Colquhoun, H., Levac, D., . . . Straus, S.E. (2018). PRISMA extension for scoping reviews (PRISMA-ScR): Checklist and explanation. *Annals of Internal Medicine*, 169(7), 467–473. <https://doi.org/10.7326/M18-0850>
- Velghe, A., De Buyser, S., Noens, L., Demuyne, R., & Petrovic, M. (2016). Hand grip strength as a screening tool for frailty in older patients with haematological malignancies. *Acta Clinica Belgica*, 71(4), 227–230. <https://doi.org/10.1080/17843286.2016.1162381>
- Walston, J., Buta, B., & Xue, Q.-L. (2018). Frailty screening and interventions: Considerations for clinical practice. *Clinics in Geriatric Medicine*, 34(1), 25–38. <https://doi.org/10.1016/j.cger.2017.09.004>
- Wilson, C.L., Chemaitilly, W., Jones, K.E., Kaste, S.C., Srivastava, D.K., Ojha, R.P., . . . Ness, K.K. (2016). Modifiable factors associated with aging phenotypes among adult survivors of childhood acute lymphoblastic leukemia. *Journal of Clinical Oncology*, 34(21), 2509–2515. <https://doi.org/10.1200/JCO.2015.64.9525>
- Wiskemann, J., Dreger, P., Schwerdtfeger, R., Bondong, A., Huber, G., Kleindienst, N., . . . Bohus, M. (2011). Effects of a partly self-administered exercise program before, during, and after allogeneic stem cell transplantation. *Blood*, 117(9), 2604–2613. <https://doi.org/10.1182/blood-2010-09-306308>
- Zhong, Y.-P., Zhang, Y.-Z., Liao, A.-J., Li, S.-X., Tian, C., & Lu, J. (2017). Geriatric assessment to predict survival and risk of serious adverse events in elderly newly diagnosed multiple myeloma patients: A multicenter study in China. *Chinese Medical Journal*, 130(2), 130–134.